

Exhibit 18



US006730502B2

(12) **United States Patent**
Van Hijum et al.

(10) **Patent No.:** **US 6,730,502 B2**
(45) **Date of Patent:** **May 4, 2004**

(54) **FRUCTOSYLTRANSFERASES**

(75) **Inventors:** **Sacha Adrianus Fokke Taco Van Hijum**, Groningen (NL); **Gerritdina Hendrika Van Geel-Schutten**, Driebergen-Rijsenberg (NL); **Lubbert Dijkhuizen**, Zuidlaren (NL); **Hakim Rahaoui**, Amersfoort (NL)

(73) **Assignee:** **Nederlandse Organisatie voor Toegepast - Natuurwetenschappelijk Onderzoek TNO**, Delft (NL)

(*) **Notice:** Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) **Appl. No.:** **09/995,587**

(22) **Filed:** **Nov. 29, 2001**

(65) **Prior Publication Data**

US 2002/0127681 A1 Sep. 12, 2002

Related U.S. Application Data

(63) Continuation-in-part of application No. 09/604,958, filed on Jun. 28, 2000, now Pat. No. 6,635,460.

(30) **Foreign Application Priority Data**

May 25, 2000 (EP) 00201872

(51) **Int. Cl.⁷** C12N 9/10; C12P 19/18;
C12P 19/04

(52) **U.S. Cl.** 435/97; 435/101; 435/193;
435/252.9

(58) **Field of Search** 435/97, 101, 193,
435/252.9

(56) **References Cited**

PUBLICATIONS

G.H. van Geel-Schutten et al. "Biochemical and Structural Characterization of the Glucan and Fructan Exopolysaccharides Synthesized by the *Lactobacillus reuteri* Wild-Type Strain and by Mutant Strains", *Applied Environ. Microbiol.* 65(7); 3008-3014. (Jul. 1).*

G.H. van Geel-Schutten et al. "Screening and Characterization of *Lactobacillus* Strains Producing Large Amounts of Exopolysaccharides", *Appl. Microbiol. Biotechnol.* 50: 697-703. (1998).*

* cited by examiner

Primary Examiner—Rebecca Prouty

(74) *Attorney, Agent, or Firm*—Young & Thompson

(57) **ABSTRACT**

The present invention describes two novel proteins having fructosyltransferase activity. Both enzymes are derived from lactobacilli, which are food-grade micro-organisms with the Generally Recognized As Safe (GRAS) status. One of these proteins produces an inulin and fructo-oligosaccharides, while the other produces a levan and fructo-oligosaccharides. According to the invention lactobacilli capable of producing an inulin and/or a levan and/or fructo-oligosaccharides using one or both of the fructosyltransferases can be used as a probiotic or a symbiotic.

8 Claims, 8 Drawing Sheets

Fig 1 (1)

1 ~~tacnattggg~~ ~~tggcgggggt~~ ~~gaagaaacgg~~ ggttacttct atgctagaac gcaaggaaaca 19ft>

y n g v a e v k k r g y f y a r t

y n g v a e v n t e r q a n g q i

61 taaaaaaatg tataaaagcg gtaaaaattg ggcagtcggt acactctcga ctgctgcgct

1 m y k s g k n v a v v t l s t a a

121 ggtatttggt gcaacaactg taaatgcac cyccggacaca aatattgaaa acaatgatcc

18 l v f g a t t v n a s a d t n i e n n d

181 ttctactgta caagttacaa caggtgataa tgatattgct gttaaaagtg tgacacttgg

38 s s t v q v t t g d n d i a v k s v t l

241 tagtgggtcaa gttagtgcag ctagtgtac gactattaga acttctgcta atgcaaatag

58 g s g q v s a a s d t t i r t s a n a n

301 tgctttcttct gccgctaata cacaaaattc taacagtcac gtagcaagtt ctgctgcaat

78 s a s s a a n t q n s n s q v a s s a a

361 aacatcatct acnagttccg cagcttcatt aaataacaca gatagtaaag cggctcaaga

98 i t s s t s s a a s l n n t d s k a a q

421 aaatactaata acagccaaaa atgatgacac gcaaaaagct gcaccagcta acgaatcttc

118 e n t n t a k n d d t q k a a p a n e s

481 tgaagctaaa aatgaaccag ctgtaaaagt taatgattct tcagctgcaa aaaatgatga

138 s e a k n e p a v n v n d s s a a k n d

541 tcaacaatcc agtaaaaaga atactaccgc taagttaaac aaggatgctg aaaacgttgt

158 d q q s s k k n t t a k l n k d a e n v

601 aaaaaaggcg ggaattgatc etaacagttt aactgatgac cagattaaag cattaaataa

178 v k k a g i d p n s l t d d q i k a l n

Fig 1 (2)

661 gatgaacttc tcgaaagctg caaagtctgg tacacaaatg acttataatg atttcaaaa
198 k m n f s k a a k s g t q m t y n d f q

721 gattgctgat acgttaatca aagaagatgg tgggtacaca gtccattct ttaaagcaag 20ftfi <
218 k i a d t l i k q d g r y t v p f f k a

781 tgaatcaaaa aatargcctg ccgtacaac taaagatgca caaactaata ctattgaacc
238 s e i k n m p a a t t k d a q t n t i e

841 tttagatgta tgggattcat ggccagtcca agatgttcgg acaggacaag ttgctaattg 5ftf >
258 p l d v w d s w p v q d v r t g q v a n 8ftfi <

901 gaatggctat caacttgcca tcgcaatgat gggaattcca aaccaaaatg ataactat
278 w n g y q l v i a m m g i p n q n d n h

961 ctatctctta tataataagt atggtgataa tgaattaagt cattggaaga atgtagggtcc 7ftf >
298 i y l l y n k y g d n e l s h v k n v g

1021 aatttttggc tataattcta ccgcggttc agagatggg tcaggatcag ctgttttgaa 7ftf >
318 p i f g y n s t a v s q e w s g s a v l 6ftfi <

1081 cagtgaatac tctatccaat tattttatac aagggtagac acgtctgata acaataccaa
338 n s d n s i q l f y t r v d t s d n n t

1141 tcacaaaaa attgctagcg ctactcttta tttactgat antattgga atgtatcact NheI
358 n h q k i a s a t l y l t d n n g n v s AC1(i)<>

1201 cgctcaggta cgaaatgact atattgtatt tgaaggatgat ggctattact accanaactta AC2(i)<>
378 l a q v r n d y i v f e g d g y y y q t

1261 tgatcaatgg aagctacta acaaaggatgc cgataatatt gcaatgcgtg atgctcatgt
398 y d q w k a t n k g a d n i a m r d a h

Fig 1 (3)

1321 aattgaagat ggtaatggtg atcggtacct tgtttttgaa gcaagtactg gtttgaaaa
418 v i e d g n g d r y l v f e a s t g l e

1381 ttatcaaggc gaggaccaa tttataactg gttaaattat ggcggagatg acgcatttaa
438 n y q g e d q i y n v l n y g g d d a f

1441 tatcaagagc ttatttagaa ttctttccaa tgatgatatt aagagtcggg caacttgggc
458 n i k s l f r i l s n d d i k s r a t v

1501 taatgcagct atcggtatcc tcaactaaa taaggacgaa aagaatccta aggtggcaga
478 a n a a i g i l k l n k d e k n p k v a

1561 gttatactca ccattaattt ctgcaccaat ggtaagcgat gaaattgagc gaccaaattg
498 e l y s p l i s a p n v s d e i e r p n

1621 agttaaatta ggtantaat attacttatt tgcgcctacc cytttaaate gaggaagtaa
518 v v k l g n k y y l f a a t r l n r g s

1681 tgatgatgct tggatgaatg ctaattatgc cgttggtgat aatgttgcaa tggtcggata
538 n d d a w m n a n y a v g d n v a n v g

1741 tgttgctgat agtctaactg gatcttataa gccattaaat gattctggag tagtcttgac
558 y v a d s l t g s y k p l n d s g v v l

1801 tgcttctgtt cctgcaact ggcggacagc aacttattca tattatgctg tcccogttgc
578 t a s v p a n w r t a t y s y y a v p v

1861 cggaaaagat gaccaagtat tagttacttc atatatgact aatagaaatg gagtagcggg
598 a g k d d q v l v t s y m t n r n g v a

1921 taaaggaatg gattenactt ggcacagag tttcttacta caaattaacc cggataaac 12ftfi <
618 g k g m d s t v a p s f l l q i n p d n

Fig 1 (4)

1981 aactactggt ttagctaaaa tgactaatca aggggattgg atttgggatg attcaagcga

638 t t t v l a k m t n q g d w i w d d s s

2041 aaatcttgat atgattggtg atttagactc cgctgcttta cctggcgaac gtgataaacc

658 e n l d m i g d l d s a a l p g e r d k

2101 tgttgattgg gacttaattg gttatggatt aaaaccgcat gatcctgcta caccaaatga

678 p v d w d l i g y g l k p h d p a t p n

2161 tcctgaaacg ccaactacac cagaaacccc tgagacacct aatactccca aaacaccaa

698 d p e t p t t p e t p e t p n t p k t p

2221 gactcctgaa aatcctggga cacctcaaac tcctaataca cctaatactc cggaaattcc

718 k t p e n p g t p q t p n t p n t p e i

2281 tttaactcca gaaacgccta agcaacctga aacccaaact aataatcggt tgccacaaac

738 p l t p e t p k q p e t q t n n r l p q

2341 tgguaataat gccataaag ccattgattgg cctaggatatg ggaacattgc ttagtatggt

758 t g n n a n k a m i g l g m g t l l s m

2401 tggctcttgcg gaaattaaca aacgtcgatt taactaata ctttaaaata aaaccgctaa

778 f g l a e i n k r r f n

2461 gccttaaatt cagcttaacg gttttttatt ttaaaagttt ttattgtaaa aaagcgaatt

2521 atcattaata ctaatgcaat tgttgtaaga ccttacgaca gtagtacaa tgaatttgcc

2581 catctttgtc gg




Fig 3

The N-terminal sequence of FTFB (levansucrase):

(A)QVESNNYNGVAEVNTERQANGQI(G)(V)(D).


Internal peptide sequences of FTFB (levansucrase):

- **(M)(A)HLDVWDSWPVQDP(V),**
- **NA·GSIFGT(K),**
- **V(E)(E)VYSPKVSTLMASDEVE.**

Fig 4

5ftf			
<i>B. amyloliquefaciens</i> SacB	80	GLDVVDSWFLQAD	93
<i>B. subtilis</i> SacB	82	GLDVVDSWFLQAD	95
<i>S. mutans</i> SacB	243	DLDVVDSWFLVQDAK	256
<i>S. salivarius</i> Ftf	282	ELDVVDSWFLVQDAK	295
		:*****:*:*.	
6ftf			
<i>B. amyloliquefaciens</i> SacB	156	QTQENSGSATFTSDGK	171
<i>B. subtilis</i> SacB	158	QTQENSGSATFTSDGK	173
<i>S. mutans</i> SacB	312	LTQENSGSATVNEDEGS	327
<i>S. salivarius</i> Ftf	351	DDQQNSGSATVNSDGS	366
		*:*****...**.	
12ftf			
<i>B. amyloliquefaciens</i> SacB	440	KATFGPSFLMN	450
<i>B. subtilis</i> SacB	440	QSTFAPSFLIN	450
<i>S. mutans</i> SacB	609	NSTWAPSFLIQ	619
<i>S. salivarius</i> Ftf	655	KSTWAPSFLIK	665
		::*:.****:::	

Fig 5



A

PCR

B

PCR

C

PCR

D

Inverse PCR

E

PCR

Inverse PCR

1

FRUCTOSYLTRANSFERASES

CROSS-REFERENCE TO RELATED APPLICATION

This application is a continuation-in-part of application of U.S. application Ser. No. 09/604,958 filed on Jun. 28, 2000, now U.S. Pat. No. 6,635,460, which claims priority from European Application No. 00201872.9 filed on May 25, 2000.

The present invention is in the field of enzymatic production of biomolecules. The invention is particularly concerned with two novel fructosyltransferases derived from lactobacilli and with a process for recombinant production of the enzymes and for the production of useful levans, inulins and fructo-oligosaccharides from sucrose.

BACKGROUND OF THE INVENTION

Lactic acid bacteria (LAB) play an important role in the fermentative production of food and feed. Traditionally, these bacteria have been used for the production of for instance wine, beer, bread, cheese and yoghurt, and for the preservation of food and feed, e.g. olives, pickles, sausages, sauerkraut and silage. Because of these traditional applications, lactic acid bacteria are food-grade microorganisms that possess the Generally Recognised As Safe (GRAS) status. Due to the different products which are formed during fermentation with lactic acid bacteria, these bacteria contribute positively to the taste, smell and preservation of the final product. The group of lactic acid bacteria encloses several genera such as *Lactobacillus*, *Leuconostoc*, *Pediococcus*, *Streptococcus*, etc.

In recent years also the health promoting properties of lactic acid bacteria have received much attention. They produce an abundant variety of exopolysaccharides (EPS's). These polysaccharides are thought to contribute to human health by acting as prebiotic substrates, nutraceuticals, cholesterol lowering agents or immunomodulators.

To date high molecular weight polysaccharides produced by plants (such as cellulose, starch and pectin), seaweeds (such as alginate and carrageenan) and bacteria (such as alginate, gellan and xanthan) are used in several industrial applications as viscosifying, stabilising, emulsifying, gelling or water binding agents. Although all these polysaccharides are used as food additives, they originate from organisms not having the GRAS status. Thus they are less desirable than the exopolysaccharides of microorganisms, such as lactic acid bacteria, which have the GRAS status.

The exopolysaccharides produced by LAB can be divided in two groups, heteropolysaccharides and homopolysaccharides; these are synthesized by totally different mechanisms. The former consist of repeating units in which residues of different types of sugars are present and the latter consist of one type of monosaccharide. The synthesis of heteropolysaccharides by lactic acid bacteria, including lactobacilli, has been studied extensively in recent years. Considerably less information is available on the synthesis of homopolysaccharides from lactobacilli, although some studies have been performed. Homopolysaccharides with fructose as the constituent sugar can be divided into two groups, inulins and levans. Inulins consist of 2,1-linked β -fructofuranoside residues, whereas levans consist of 2,6-linked β -fructofuranoside residues. Both can be linear or branched. The size of bacterial levans can vary from 20 kDa up to several MDa. There is limited information on the synthesis of levans. In most detail this synthesis has been studied in *Zymomonas mobilis* and in *Bacillus* species.

2

Within lactic acid bacteria, fructosyltransferases have only been studied in streptococci. So far no fructosyltransferases have been reported in lactobacilli.

In a recent report the *Lactobacillus reuteri* strain LB 121 was found to produce both a glucan and a fructan when grown on sucrose, but only a fructan when grown on raffinose (van Geel-Schutten, G. H. et al., Appl. Microbiol. Biotechnol. (1998) 50, 697-703). In another report the glucan and fructan were characterised by their molecular weights (of 3,500 and 150 kDa respectively) and the glucan was reported to be highly branched with a unique structure consisting of a terminal, 4-substituted, 6-substituted, and 4,6-disubstituted α -glucose in a molar ratio 1.1: 2.7:1.5:1.0 (van Geel-Schutten, G. H. et al., Appl. Environ. Microbiol. (1999) 65, 3008-3014). The fructan was identified as a linear (2 \rightarrow 6)- β -D-fructofuranan (also called a levan). This was the first example of fructan synthesis by a *Lactobacillus* species.

SUMMARY OF THE INVENTION

Two novel genes encoding enzymes having fructosyltransferase activity have now been found in *Lactobacillus reuteri*, and their amino acid sequences have been determined. These are the first two enzymes identified in a *Lactobacillus* species capable of producing a fructan. One of the enzymes is an inulosucrase which produces a high molecular weight (>10⁷ Da) fructan containing β (2-1) linked fructosyl units and fructo-oligosaccharides, while the other is a levansucrase which produces a fructan containing β (2-6) linked fructosyl units. The invention thus pertains to the enzymes, to DNA encoding them, to recombinant cells containing such DNA and to their use in producing carbohydrates, as defined in the appending claims.

DESCRIPTION OF THE INVENTION

It was found according to the invention that one of the novel fructosyltransferases (FTFA; an inulosucrase) produces a high molecular weight inulin with β (2-1) linked fructosyl units and fructo-oligosaccharides. The fructo-oligosaccharides synthesis was also observed in certain *Lactobacillus* strains, in particular in certain strains of *Lactobacillus reuteri*. However, the inulin has not been found in *Lactobacillus reuteri* culture supernatants, but only in extracts of *E. coli* cells expressing the above-mentioned fructosyltransferase. This inulosucrase consists of either 798 amino acids (2394 nucleotides) or 789 amino acids (2367 nucleotides) depending on the potential start codon used. The molecular weight (MW) deduced of the amino acid sequence of the latter form is 86 kDa and its isoelectric point is 4.51, at pH 7.

The amino acid sequence of the inulosucrase is shown in SEQ ID No. 1 (FIG. 1, amino acid residues 1-789). As mentioned above, the nucleotide sequence contains two putative start codons leading to either a 2394 (see SEQ ID No. 3) or 2367 (see SEQ ID No. 2) nucleotide form of the inulosucrase. Both putative start codons are preceded by a putative ribosome binding site, GGGG (located 12 base pairs upstream its start codon) or AGGA (located 14 base pairs upstream its start codon), respectively (see FIG. 1 and SEQ ID No. 4).

The present invention covers a protein having inulosucrase activity with an amino acid identity of at least 65%, preferably at least 75%, and more preferably at least 85%, compared to the amino acid sequence of SEQ ID No. 1. The invention also covers a part of a protein with at least 15 contiguous amino acids which are identical to the corresponding part of the amino acid sequence of SEQ ID No. 1.

Fructosyltransferases have been found in several bacteria such as *Zymomonas mobilis*, *Erwinia amylovora*, *Acetobacter amylovora*, *Bacillus polymyxa*, *Bacillus amyloliquefaciens*, *Bacillus stearothermophilus*, and *Bacillus subtilis*. In lactic acid bacteria this type of enzyme previously has only been found in some streptococci. Most bacterial fructosyltransferases have a molecular mass of 50–100 kDa (with the exception of the fructosyltransferase found in *Streptococcus salivarius* which has a molecular mass of 140 kDa). Amino acid sequence alignment revealed that the novel inulosucrase of lactobacilli has high homology with fructosyltransferases originating from Gram positive bacteria, in particular with Streptococcus enzymes. The highest homology (FIG. 2) was found with the SacB enzyme of *Streptococcus mutans* Ingbritt A (62% identity within 539 amino acids).

Certain putative functions based on the alignment and site-directed mutagenesis studies can be ascribed to several amino acids of the novel inulosucrase. Asp-263, Glu-330, Asp-415, Glu-431, Asp-511, Glu-514, Arg-532 and/or Asp-551 of the amino acid sequence of SEQ ID No. 1 are identified as putative catalytic residues. Noteworthy, a hydrophobicity plot according to Kyte and Doolittle (1982) J. Mol. Biol. 157, 105–132 suggests that the novel inulosucrase contains a putative signal sequence according to the Von Heijne rule. The putative signal peptidase site is located between Gly at position 21 and Ala at position 22. Furthermore, it is striking that the C-terminal amino acid sequence of the novel inulosucrase contains a putative cell wall anchor amino acid signal LPXTG (SEQ ID No. 5) and a 20-fold repeat of the motif PXX (residues 690–749 of SEQ ID NO: 1) (see figure 1.), where P is proline and X is any other amino acid. In 15 out of 20 repeats, however, the motif is PXT. This motif has so far not been reported in proteins of prokaryotic and eukaryotic origin.

A nucleotide sequence encoding any of the above mentioned proteins, mutants, variants or parts thereof is also a subject of the invention. Furthermore, the nucleic acid sequences corresponding to expression-regulating regions (promoters, enhancers, terminators) of at least 30 contiguous nucleic acids contained in the nucleic acid sequence (-67)-(-1) or 2367–2525 of SEQ ID No. 4 (see also FIG. 1) can be used for homologous or heterologous expression of genes. Such expression-regulating sequences are operationally linked to a polypeptide-encoding nucleic acid sequence such as the genes of the fructosyltransferase according to the invention. A nucleic acid construct comprising the nucleotide sequence operationally linked to an expression-regulating nucleic acid sequence is also covered by the invention.

A recombinant host cell, such as a mammalian (with the exception of human), plant, animal, fungal or bacterial cell, containing one or more copies of the nucleic acid construct mentioned above is an additional subject of the invention. The inulosucrase gene (starting at nucleotide 41) has been cloned in an *E. coli* expression vector under the control of an ara promoter in *E. coli* Top10. *E. coli* Top10 cells expressing the recombinant inulosucrase hydrolysed sucrose and synthesized fructan material. SDS-PAGE of arabinose induced *E. coli* Top10 cell extracts suggested that the recombinant inulosucrase has a molecular weight of 80–100 kDa, which is in the range of other known fructosyltransferases and in line with the molecular weight of 86 kDa deduced of the amino acid sequence depicted in FIG. 1.

The invention further covers an inulosucrase according to the invention which, in the presence of sucrose, produces a inulin having $\beta(2-1)$ -linked D-fructosyl units and fructo-

oligosaccharides. Two different types of fructans, inulins and levans, exist in nature. Surprisingly, the novel inulosucrase expressed in *E. coli* Top10 cell synthesizes a high molecular weight ($>10^7$ Da) inulin and fructo-oligosaccharides, while in *Lactobacillus reuteri* culture supernatants, in addition to the fructo-oligosaccharides, a levan and not an inulin is found. This discrepancy can have several explanations: the inulosucrase gene may be silent in *Lactobacillus reuteri*, or may not be expressed in *Lactobacillus reuteri* under the conditions tested, or the inulosucrase may only synthesize fructo-oligosaccharides in its natural host, or the inulin polymer may be degraded shortly after synthesis, or may not be secreted and remains cell-associated, or the inulosucrase may have different activities in *Lactobacillus reuteri* and *E. coli* Top10 cells.

It was furthermore found according to the invention that certain lactobacilli, in particular *Lactobacillus reuteri*, possess another fructosyltransferase, a levansucrase (FTFB), in addition to the inulosucrase described above. The N-terminal amino acid sequence of the fructosyltransferase purified from *Lactobacillus reuteri* supernatant was found to be QVESNNYNGVAEVNTERQANGQI (residues 2–24 of SEQ ID No. 6). Furthermore, three internal sequences were identified, namely (M)(A)HLDVWDSWPVQDP(V) (SEQ ID No. 7), NAGSIFGT(K) (SEQ ID No. 8), V(E) (E) VYSPKVSTLMASDEVE (SEQ ID No. 9). The N-terminal amino acid sequence could not be identified in the deduced inulosucrase sequence. Also the amino acid sequences of the three internal peptide fragments of the purified fructosyltransferase were not present in the putative inulosucrase sequence. Evidently, the inulosucrase gene does not encode the purified fructosyltransferase synthesizing the levan. The complete amino acid sequence of the levansucrase is shown in SEQ ID No. 11 and the nucleotide sequence is shown in SEQ ID No. 10. The levansucrase comprises a putative membrane anchor (see amino acids 761–765 in SEQ ID No. 11) and a putative membrane spanning domain (see amino acids 766–787 in SEQ ID No. 11). The fructan produced by the levansucrase was identified in the *Lactobacillus reuteri* culture supernatant as a linear (2 \rightarrow 6)- β -D-fructofuranan with a molecular weight of 150 kDa. The purified enzyme also produces this fructan.

Additionally, the invention thus covers a protein having levansucrase activity with an amino acid identity of at least 65%, preferably at least 75%, and more preferably at least 85%, compared to the amino acid sequence of SEQ ID NO. 11. The second novel fructosyltransferase produces a high molecular weight fructan with $\beta(2-6)$ linked fructosyl units with sucrose or raffinose as substrate. The invention also covers a part of a protein with least 15 contiguous amino acids, which are identical to the corresponding part of the amino acid sequence of SEQ ID No. 11. A nucleotide sequence encoding any of the above-mentioned proteins, mutants, variants or parts thereof is a subject of the invention as well as a nucleic acid construct comprising the nucleotide sequence mentioned above operationally linked to an expression-regulating nucleic acid sequence. A recombinant host cell, such as a mammalian (with the exception of human), plant, animal, fungal or bacterial cell, containing one or more copies of the nucleic acid construct mentioned above is an additional subject of the invention. The invention further covers a protein according to the invention which, in the presence of sucrose, produces a fructan having $\beta(2-6)$ -linked D-fructosyl units.

The invention also pertains to a process of producing an inulin-type and/or a levan-type of fructan as described above using fructosyltransferases according to the invention and a

suitable fructose source such as sucrose, stachyose or raffinose. The fructans may either be produced by *Lactobacillus* strains or recombinant host cells according to the invention containing one or both fructosyl transferases or by a fructosyltransferase enzyme isolated by conventional means from the culture of fructosyltransferase-positive *Lactobacilli*, especially a *Lactobacillus reuteri*, or from a recombinant organism containing the fructosyltransferase gene or genes.

Additionally, the invention concerns a process of producing fructo-oligosaccharides containing the characteristic structure of the fructans described above using a *Lactobacillus* strain or a recombinant host cell according to the invention containing one or both fructosyltransferases or an isolated fructosyltransferase according to the invention. There is a growing interest in oligosaccharides derived from homopolysaccharides, for instance for prebiotic purposes. Several fructo- and gluco-oligosaccharides are known to stimulate the growth of bifidobacteria in the human colon. Fructo-oligosaccharides produced by the fructosyltransferase described above are also part of the invention. Another way of producing fructo-oligosaccharides is by hydrolysis of the fructans described above. This hydrolysis can be performed by known hydrolysis methods such as enzymatic hydrolysis with enzymes such as levansase or inulinase or by acid hydrolysis. The fructo-oligosaccharides can also be produced in the presence of a fructosyltransferase according to the invention and an acceptor molecule such as lactose or maltose. The fructo-oligosaccharides to be produced according to the invention preferably contain at least 2, more preferably at least 3, up to about 20 anhydrofructose units, optionally in addition to one or more other (glucose, galactose, etc.) units. These fructo-oligosaccharides are useful as prebiotics, and can be administered to a mammal in need of improving the bacterial status of the colon.

The invention also concerns chemically modified fructans and fructo-oligosaccharides based on the fructans described above. Chemical modification can be achieved by oxidation, such as hypochlorite oxidation resulting in ring-opened 2,3-dicarboxy-anhydrofructose units (see e.g. EP-A-427349), periodate oxidation resulting in ring-opened 3,4-dialdehyde-anhydrofructose units (see e.g. WO 95/12619), which can be further oxidised to (partly) carboxylated units (see e.g. WO 00/26257), TEMPO-mediated oxidation resulting in 1- or 6-carboxy-anhydrofructose units (see e.g. WO 95/07303). The oxidised fructans have improved water-solubility, altered viscosity and a retarded fermentability and can be used as metal-complexing agents, detergent additives, strengthening additives, bioactive carbohydrates, emulsifiers and water binding agents. They can also be used as starting materials for further derivatisation such as cross-linking and the introduction of hydrophobes. Oxidised fructans coupled to amino compounds such as proteins, or fatty acids can be used as emulsifiers and stabilizers. (Partial) hydrolysis of fructans according to the invention and modified fructans according to the invention results in fructo-oligosaccharides, which can be used as bioactive carbohydrates or prebiotics. The oxidised fructans of the invention preferably contain 0.05–1.0 carboxyl groups per anhydrofructose unit, e.g. as 6- or 1-carboxyl units.

Another type of chemical modification is phosphorylation, as described in O.B. Wurzburg (1986) *Modified Starches: properties and uses*. CRC Press Inc., Boca Raton, 97–112. One way to achieve this modification is by dry heating fructans with a mixture of monosodium and disodium hydrogen phosphate or with tripolyphosphate. The phosphorylated fructans are suitable as wet-end additives in

papermaking, as binders in paper coating compositions, as warp sizing-agents, and as core binders for sand molds for metal casting. A further type of derivatisation of the fructans is acylation, especially acetylation using acetic or propionic anhydride, resulting in products suitable as bleaching assistants and for the use in foils. Acylation with e.g. alkenyl succinic anhydrides or (activated) fatty acids results in surface-active products suitable as e.g. surfactants, emulsifiers, and stabilizers.

Hydroxyalkylation, carboxymethylation, and aminoalkylation are other methods of chemical derivatisation of the fructans. Hydroxyalkylation is commonly performed by base-catalysed reaction with alkylene oxides, such as ethylene oxide, propylene oxide or epichlorohydrine; the hydroxyalkylated products have improved solubility and viscosity characteristics. Carboxymethylation is achieved by reaction of the fructans with mono-chloroacetic acid or its alkali metal salts and results in anionic polymers suitable for various purposes including crystallisation inhibitors, and metal complexants. Amino-alkylation can be achieved by reaction of the fructans with alkylene imines, haloalkyl amines or amino-alkylene oxides, or by reaction of epichlorohydrine adducts of the fructans with suitable amines. These products can be used as cationic polymers in a variety of applications, especially as a wet-end additive in paper making to increase strength, for filler and fines retention, and to improve the drainage rate of paper pulp. Other potential applications include textile sizing and wastewater purification. The above mentioned modifications can be used either separately or in combination depending on the desired product. Furthermore, the degree of chemical modification is variable and depends on the intended use. If necessary 100% modification, i.e. modification of all anhydrofructose units can be performed. However, partial modification, e.g. from 1 modified anhydrofructose unit per 100 up to higher levels, will often be sufficient in order to obtain the desired effect. The modified fructans have a DP (degree of polymerisation) of at least 100, preferably at least 1000 units.

Use of a *Lactobacillus* strain capable of producing a levan, inulin or fructo-oligosaccharides or a mixture thereof, as a probiotic, is also covered by the invention. Preferably, the *Lactobacillus* strain is also capable of producing a glucan, especially an 1,4/1,6- α -glucan as referred to above. The efficacy of some *Lactobacillus reuteri* strains as a probiotics has been demonstrated in various animals such as for instance poultry and humans. The administration of some *Lactobacillus reuteri* strains to pigs resulted in significantly lower serum total and LDL-cholesterol levels, while in children *Lactobacillus reuteri* is used as a therapeutic agent against acute diarrhea. For this and other reasons *Lactobacillus reuteri* strains, which were not reported to produce the glucans or fructans described herein, have been supplemented to commercially available probiotic products. The mode of action of *Lactobacillus reuteri* as a probiotic is still unclear. Preliminary studies indicated that gut colonization by *Lactobacillus reuteri* may be of importance. According to the invention, it was found that the mode of action of *Lactobacillus reuteri* as a probiotic may reside partly in the ability to produce polysaccharides. *Lactobacillus* strains, preferably *Lactobacillus reuteri* strains, and more preferably *Lactobacillus reuteri* strain LB 121 and other strains containing one or more fructosyltransferase genes encoding proteins capable of producing inulins, levans and/or fructo-oligosaccharides can thus advantageously be used as a probiotic. They can also, together with these polysaccharides, be used as a symbiotic (instead of the term symbiotic, the term synbiotic can also be used). In that

respect another part of the invention concerns a probiotic or symbiotic composition containing a *Lactobacillus* strain capable of producing an inulin, a levan or fructo-oligosaccharides and/or a glucan or a mixture thereof, said production being performed according to the process according to the invention. The probiotic or symbiotic compositions of the invention may be directly ingested with or without a suitable vehicle or used as an additive in conjunction with foods. They can be incorporated into a variety of foods and beverages including, but not limited to, yoghurts, ice creams, cheeses, baked products such as bread, biscuits and cakes, dairy and dairy substitute foods, confectionery products, edible oil compositions, spreads, breakfast cereals, juices and the like.

Furthermore, the invention pertains to a process of improving the microbial status in the mammalian colon comprising administering an effective amount of a *Lactobacillus* strain capable of producing an oligosaccharide or polysaccharide according to the invention and to a process of improving the microbial status of the mammalian colon comprising administering an effective amount of an oligosaccharide or polysaccharide produced according to the process according to the invention.

EXAMPLES

EXAMPLE 1

Isolation of DNA from *Lactobacillus reuteri* Nucleotide Sequence Analysis of the Inulosucrase (fifA) Gene, Construction of Plasmids for Expression of the Inulosucrase Gene in *E. coli* Top10 Expression of the Inulosucrase gene in *E. coli* Top10 and Identification of the Produced Polysaccharides Produced by the Recombinant Enzyme.

General procedures for cloning, DNA manipulations and agarose gel electrophoresis were essentially as described by Sambrook et al. (1989) Molecular cloning: a laboratory manual, 2nd ed. Cold Spring Harbour Laboratory Press, Cold Spring Harbour, N.Y. Restriction endonuclease digestions and ligations with T4 DNA ligase were performed as recommended by the suppliers. DNA was amplified by PCR techniques using ampliTAQ DNA polymerase (Perkin Elmer) or Pwo DNA polymerase. DNA fragments were isolated from agarose gels using the Qiagen extraction kit (Qiagen GmbH), following the instructions of the suppliers. *Lactobacillus reuteri* strain 121 (LMG 18388) was grown at 37° C. in MRS medium (DIFCO) or in MRS-s medium (MRS medium containing 100 g/l sucrose instead of 20 g/l glucose). When fructo-oligosaccharides production was investigated phosphate was omitted and ammonium citrate was replaced by ammonium nitrate in the MRS-s medium. *E. coli* strains were grown aerobically at 37° C. in LB medium, where appropriate supplemented with 50 µg/ml ampicillin (for selection of recombinant plasmids) or with 0.02% (w/v) arabinose (for induction of the inulosucrase gene).

Total DNA of *Lactobacillus reuteri* was isolated according to Verhasselt et al. (1989) FEMS Microbiol. Lett. 59, 135-140 as modified by Nagy et al. (1995) J. Bacteriol. 177, 676-687.

The inulosucrase gene was identified by amplification of chromosomal DNA of *Lactobacillus reuteri* with PCR using degenerated primers (5 fif, 6 ftfi, and 12 ftfi, see table 1) based on conserved amino acid sequences deduced from different bacterial fructosyltransferase genes (SacB of *Bacillus amyloliquefaciens*, SacB of *Bacillus subtilis*, *Streptococcus mutans* fructosyltransferase and *Streptococcus salivarius* fructosyltransferase, see FIG. 4) and *Lactobacillus*

reuteri DNA as template. Using primers 5 fif and 6 ftfi, an amplification product with the predicted size of about 234 bp was obtained (FIG. 5A). This 234 bp fragment was cloned in *E. coli* JM109 using the pCR2.1 vector and sequenced. Transformations were performed by electroporation using the BioRad gene pulser apparatus at 2.5 kV, 25 µF and 200 Ω, following the instructions of the manufacturer. Sequencing was performed according to the method of Sanger et al. (1977) Proc. Natl. Acad. Sci. USA 74, 5463-5467. Analysis of the obtained sequence data confirmed that part of a fructosyltransferase (fif gene had been isolated. The 234 bp amplified fragment was used to design primers 7 fif and 8 ftfi (see table 1). PCR with the primers 7 fif and 12 ftfi gave a product of the predicted size of 948 bp (see FIG. 5B); its sequence showed clear similarity with previously characterized fructosyltransferase genes. The 948 bp amplified fragment was used to design the primers fifAC1(i) and fifAC2(i) (see table 1) for inverse PCR. Using inverse PCR techniques a 1438 bp fragment of the inulosucrase gene was generated, including the 3' end of the inulosucrase gene (see FIG. 5C). The remaining 5' fragment of the inulosucrase gene was isolated with a combination of standard and inverse PCR techniques. Briefly, *Lactobacillus reuteri* DNA was cut with restriction enzyme XhoI and ligated. PCR with the primers 7 fif and 8 ftfi, using the ligation product as a template, yielded a 290 bp PCR product which was cloned into pCR2.1 and sequenced. This revealed that primer 8 ftfi had annealed aspecifically as well as specifically yielding the 290 bp product (see FIG. 5D).

At this time, the N-terminal amino acid sequence of a fructosyltransferase enzyme (FTFB) purified from the *Lactobacillus reuteri* strain 121 was obtained. This sequence consisted of the following 23 amino acids: QVESNNYN-GVAEVNTERQANGQI (residues 2-24 of SEQ ID No. 6). The degenerated primer 19 fif (YNGVAEV) (residues 8-14 of SEQ ID NO: 6) was designed on the basis of a part of this N-terminal peptide sequence and primer 20 ftfi was designed on the 290 bp PCR product. PCR with primers 19 fif and 20 ftfi gave a 754 bp PCR product (see FIG. 5E), which was cloned into pCR2.1 and sequenced. Both DNA strands of the entire fructosyltransferase gene were double sequenced. In this way the sequence of a 2.6 kb region of the *Lactobacillus reuteri* DNA, containing the inulosucrase gene and its surroundings were obtained.

The plasmids for expression of the inulosucrase gene in *E. coli* Top10 were constructed as described hereafter. A 2414 bp fragment, containing the inulosucrase gene starting at the first putative start codon at position 41, was generated by PCR, using primers fifA1 and fifA2i. Both primers contained suitable restriction enzyme recognition sites (a NcoI site at the 5' end of fifA1 and a BglII site at the 3' end of fifA2i). PCR with *Lactobacillus reuteri* DNA, Pwo DNA polymerase and primers fifA1 and fifA2i yielded the complete inulosucrase gene flanked by NcoI and BglII restriction sites. The PCR product with blunt ends was ligated directly into pCRbluntII-Topo. Using the NcoI and BglII restriction sites, the putative fifA gene was cloned into the expression vector pBAD, downstream of the inducible arabinose promoter and in frame upstream of the Myc epitope and the His tag. The pBAD vector containing the inulosucrase gene (pSVH101) was transformed to *E. coli* Top10 and used to study inulosucrase expression. Correct construction of plasmid containing the complete inulosucrase gene was confirmed by restriction enzyme digestion analysis and by sequence analysis, showing an in frame cloning of the inulosucrase gene using the ribosomal binding site provided by the pBAD vector and the first putative start codon (at position 41) of inulosucrase (see FIG. 1).

Plasmid DNA of *E. coli* was isolated using the alkaline lysis method of Birnboim and Doly (1979) Nucleic Acids Res. 7, 1513–1523 or with a Qiagen plasmid kit following the instructions of the supplier. Cells of *E. coli* Top10 with pSVH101 were grown overnight in LB medium containing 0.02% (w/v) arabinose and were harvested by centrifugation. The pellet was washed with 25 mM sodium acetate buffer pH 5.4 and the suspension was centrifuged again. Pelleted cells were resuspended in 25 mM sodium acetate buffer pH 5.4. Cells were broken by sonication. Cell debris and intact cells were removed by centrifugation for 30 min at 4° C. at 10,000×g and the resulting cell free extract was used in the enzyme assays.

The fructosyltransferase activities were determined at 37° C. in reaction buffer (25 mM sodium acetate, pH 5.4, 1 mM CaCl₂, 100 g/l sucrose) by monitoring the release of glucose from sucrose, by detecting fructo-oligosaccharides or by determining the amount of fructan polymer produced using *E. coli* cell free extracts or *Lactobacillus reuteri* culture supernatant as enzyme source. Sucrose, glucose and fructose were determined enzymatically using commercially available kits.

Fructan production by *Lactobacillus reuteri* was studied with cells grown in MRS-s medium. Product formation was also studied with cell-free extracts of *E. coli* containing the novel inulosucrase incubated in reaction buffer (1 mg protein/10 ml buffer, incubated overnight at 37° C.). Fructans were collected by precipitation with ethanol. ¹H-NMR spectroscopy and methylation analysis were performed as described by van Geel-Schutten et al. (1999) Appl. Environ. Microbiol. 65, 3008–3014. The molecular weights of the fructans were determined by high performance size exclusion chromatography coupled on-line with a multi angle laser light scattering and a differential refractive index detector. Fructo-oligosaccharide synthesis was studied in *Lactobacillus reuteri* culture supernatants and in extracts of *E. coli* cells containing the novel inulosucrase incubated in reaction buffer (1 mg protein/10 ml buffer, incubated overnight at 37° C.). Glucose and fructose were determined enzymatically as described above and fructo-oligosaccharides produced were analyzed using a Dionex column. The incubation mixtures were centrifuged for 30 min at 10,000×g and diluted 1:5 in a 100% DMSO solution prior to injection on a Dionex column. A digest of inulin (DPI-20) was used as a standard. Separation of compounds was achieved with anion-exchange chromatography on a CarboPac Pa1 column (Dionex) coupled to a CarboPac PA1 guard column (Dionex). Using a Dionex GP50 pump the following gradient was generated: % eluent B is 5% (0 min); 35% (10 min); 45% (20 min); 65% (50 min); 100% (54–60 min); 5% (61–65 min). Eluent A was 0.1 M NaOH and eluent B was 0.6 M NaAc in a 0.1 M NaOH solution. Compounds were detected using a Dionex ED40 electrochemical detector with an AU working electrode and a Ag/AgCl reference-electrode with a sensitivity of 300 nC. The pulse program used was: +0.1 Volt (0–0.4 s); +0.7 Volt (0.41–0.60 s); –0.1 Volt (0.61–1.00 s). Data were integrated using a Perkin Elmer Turbochrom data integration system. A different separation of compounds was done on a cation exchange column in the calcium form (Benson BCX4). As mobile phase Ca-EDTA in water (100 ppm) was used. The elution speed was 0.4 ml/min at a column temperature of 85° C. Detection of compounds was done by a refractive index (Jasco 830-RI) at 40° C. Quantification of compounds was achieved by using the software program Turbochrom (Perkin Elmer).

SDS-PAGE was performed according to Laemmli (1970) Nature 227, 680–685 using 7.5% polyacrylamide gels. After

electrophoresis gels were stained with Coomassie Brilliant Blue or an activity staining (Periodic Acid Schiff, PAS) was carried out as described by Van Geel-Schutten et al. (1999) Appl. Environ. Microbiol. 65, 3008–3014.

TABLE 1

Nucleotide sequence of primers used in PCR reactions to identify the inulosucrase gene.

Primer name	Location (bp)	Nucleotide sequence (and SEQ ID No)
ftfAC1	1176	CTG-ATA-ATA-ATG-GAA-ATG-TAT-CAC (SEQ ID No. 12)
ftfAC2i	1243	CAT-GAT-CAT-AAG-TTT-GGT-AGT-AAT-AG (SEQ ID No. 13)
ftfAC1	1176	GTG-ATA-CAT-TTC-CAT-TAT-TAT-CAG (SEQ ID No. 14)
ftfAC2	1243	CTA-TTA-CTA-CCA-AAC-TTA-TGA-TCA-TG (SEQ ID No. 15)
ftfA1		CCA-TGG-CCA-TGG-TAG-AAC-GCA-AGG-AAC-ATA-AAA-AAA-TG (SEQ ID No. 16)
ftfA2i		AGA-TCT-AGA-TCT-GTT-AAA-TCG-ACG-TTT-GTT-AAT-TTC-TG (SEQ ID No. 17)
5ftf	845	GAY-GTN-TGG-GAY-WSN-TGG-GCC (SEQ ID No. 18)
6ftf	1052	GTN-GCN-SWN-CCN-SWC-CAY-TSY-TG (SEQ ID No. 19)
7ftf	1009	GAA-TGT-AGG-TCC-AAT-TTT-TGG-C (SEQ ID No. 20)
8ftf	864	CCT-GTC-CGA-ACA-TCT-TGA-ACT-G (SEQ ID No. 21)
12ftf	1934	ARR-AAN-SWN-GGN-GCV-MAN-GTN-SW (SEQ ID No. 22)
19ftf	1	TAY-AAY-GGN-GTN-GCN-GAR-GTN-AA (SEQ ID No. 23)
20ftf	733	CCG-ACC-ATC-TTG-TTT-GAT-TAA-C (SEQ ID No. 24)

Listed from left to right are: primer name (i, inverse primer), location (in bp) in ftfA and the sequence from 5' to 3' according to IUB group codes (N = any base; M = A or C; R = A or G; W = A or T; S = C or G; Y = C or T; K = G or T; B = not A; D = not C; H = not G; and V = not T).

EXAMPLE 2

Purification and Amino Acid Sequencing of the Levansucrase (FTFB).

Protein Purification

Samples were taken between each step of the purification process to determine the enzyme activity (by glucose GOD-Perid method) and protein content (by Bradford analysis and acrylamide gel electrophoresis). Collected chromatography fractions were screened for glucose liberating activity (GOD-Perid method) to determine the enzyme activity.

One liter of an overnight culture of LB121 cells grown on MRS medium containing 50 grams per liter maltose was centrifuged for 15 min. at 10,000×g. The supernatant was precipitated with 1.5 liter of a saturated ammonium sulphate solution. The ammonium sulphate solution was added at a rate of 50 ml/min. under continuous stirring. The resulting 60% (w/v) ammonium sulphate solution was centrifuged for 15 min. at 10,000×g. The precipitate was resuspended in 10 ml of a sodium phosphate solution (10 mM, pH 6.0) and dialysed overnight against 10 mM sodium phosphate, pH 6.0.

A hydroxylapatite column was washed with a 10 mM sodium phosphate solution pH 6.0; the dialysed sample was loaded on the column. After eluting the column with 200 mM sodium phosphate, pH 6.0 the eluted fractions were screened for glucose releasing activity and fractions were pooled for phenyl superose (a hydrophobic interactions column) chromatography. The pooled fractions were diluted

1:1 (v:v) with 25 mM sodium acetate, 2 M ammonium sulphate, pH 5.4 and loaded on a phenyl superose column (washed with 25 mM sodium acetate, 1 M ammonium sulphate, pH 5.4). In a gradient from 25 mM sodium acetate, 1 M ammonium sulphate, pH 5.4 (A) to 25 mM sodium acetate, pH 5.4 (B) fractions were collected from 35% B to 50% B.

Pooled fractions from the phenyl superose column were loaded on a gel filtration (superdex) column and eluted by a 25 mM acetate, 0.1 M sodium chloride, pH 5.4 buffer. The superdex fractions were loaded on a washed (with 25 mM sodium acetate, pH 5.4) Mono Q column and eluted with 25 mM sodium acetate, 1 M sodium chloride, pH 5.4. The fractions containing glucose liberating activity were pooled, dialysed against 25 mM sodium acetate, pH 5.4, and stored at -20° C.

A levansucrase enzyme was purified from LB121 cultures grown on media containing maltose using ammonium sulfate precipitation and several chromatography column steps (table 2). Maltose (glucose—glucose) was chosen because both glucanucrase and levansucrase can not use maltose as substrate. LB121 will grow on media containing maltose but will not produce polysaccharide. From earlier experiments it was clear that even with harsh methods the levansucrase enzyme could not be separated from its product levan. These harsh methods included boiling the levan in a SDS solution and treating the levan with HCl and TFA. No levansucrase enzyme was commercially available for the enzymatic breakdown of levan. Only a single levansucrase was detected in maltose culture supernatants. In order to prove that the enzyme purified from maltose culture supernatant is the same enzyme which is responsible for the levan production during growth on raffinose, biochemical and biophysical tests were performed.

TABLE 2

Purification of the <i>Lactobacillus reuteri</i> LB 121 levansucrase (FTFB) enzyme					
Step	Protein (mg)	Total Activity (U)	Specific Activity (U/mg)	Purification (fold)	Yield (%)
Supernatant	128	64	0.5	1	100
Ammonium sulfate precipitation (65%)	35.2	42	1.2	2.4	65.6
Hydroxyl apatite	1.5	30.6	20.4	40.8	47.8
Phenyl superose	0.27	23	85	170	36
Gel Filtration	0.055	10	182	360	16
MonoQ	0.0255	4	176	352	6

Amino Acid Sequencing of FTFB

A 5% SDS-PAA gel was allowed to "age" overnight in order to reduce the amount of reacting chemical groups in the gel. Reaction of chemicals in the PAA gel (TEMED and ammonium persulfate) with proteins can cause some undesired effects, such as N-terminal blocking of the protein, making it more difficult to determine the protein amino acid composition. 0.1 mM thioglycolic acid (scavenger to reduce the amount of reactive groups in the PAA gel material) was added to the running buffer during electrophoresis.

In order to determine the amino acid sequence of internal peptides of protein bands running in a SDS-PAA gel, protein containing bands were cut out of the PAA gel. After frac-

tionating the protein by digestion with chymotrypsin the N-terminal amino acid sequences of the digested proteins were determined (below).

N-terminal sequencing was performed by Western blotting of the proteins from the PAA gel to an Immobilon PVDF membrane (Millipore/Waters Inc.) at 0.8 mA/cm² for 1 h. After staining the PVDF membrane with Coomassie Brilliant Blue without adding acetic acid (to reduce N-terminal blocking) and destaining with 50% methanol, the corresponding bands were cut out of the PVDF membrane for N-terminal amino acid sequence determination.

Amino acid sequence determination was performed by automated Edman degradation as described by Koningsberg and Steinman (1977) The proteins (third edition) volume 3, 1-178 (Neurath and Hill, eds.). The automated equipment for Edman degradation was an Applied Biosystems model 477A pulse-liquid sequenator described by Hewick et al. (1981), J. Biol. Chem. 15, 7990-7997 connected to a RP-HPLC unit (model 120A, Applied Biosystems) for amino acid identification.

The N-terminal sequence of the purified FTFB was determined and found to be: (A) Q V E S N N Y N G V A E V N T E R Q A N G Q I (G) (V) (D) (SEQ ID No. 6). Three internal peptide sequences of the purified FTFB were determined: (M) (A) H L D V W D S W P V Q D P (V) (SEQ ID No. 7); N A G S I F G T (K) (SEQ ID No. 8); and V (E) (E) V Y S P K V S T L M A S D E V E (SEQ ID No. 9).

The following primers were designed on the basis of the N-terminal and internal peptide fragments of FTFB. Listed from left to right are: primer name, source peptide fragment and sequence (from 5' to 3'). FTFB1+FTFB3i yields approximately a 1400 bp product in a PCR reaction. FTFB1 forward (N-terminal): AA T/C-TAT-AA T/C-GG T/C-GTT-GC G/A-T/C GA-AGT (SEQ ID No. 25); and FTFB3i reverse (Internal 3): TAC-CGN-A/T C/G N-CTA-CIT-CAA-CTT (SEQ ID No. 26). The FTFB gene was partly isolated by PCR with primers FTFB1 and FTFB3i. PCR with these primers yielded a 1385 bp amplicon, which after sequencing showed high homology to *ftfA* and *SacB* from *Streptococcus mutans*.

EXAMPLE 3

Oxidation of Levans

For TEMPO-mediated oxidation, a levan according to the invention prepared as described above (dry weight 1 g, 6.15 mmol) was resuspended in 100 ml water. Next, 2,2,6,6-tetramethylpiperidine-1-oxyl (TEMPO; 1% by weight compared to the polysaccharide (0.01 g, 0.065 mmol)) was added and resuspended in 20 min. Sodium bromide (0.75 g, 7.3 mmol) was added and the suspension was cooled down to 0° C. This reaction also proceeded without bromide. A solution of hypochlorite (6 ml, 15% solution, 12.6 mmol) was adjusted to pH 10.0 with 3M HCl and cooled to 0° C. This solution was added to the suspension of the polysaccharide and TEMPO. The course of the reaction was followed by monitoring the consumption of sodium hydroxide solution, which is equivalent to the formation of uronic acid. After 30 min, 60 ml 0.1M NaOH was consumed. This amount corresponds to the formation of 97% uronic acid. Thereafter, the solution was poured out in 96% ethanol (comprising 70% of the volume of the solution) causing the product to precipitate. The white precipitate was centrifuged, resuspended in ethanol/water (70/30 v/v) and centrifuged again. Next, the precipitate was resuspended in 96% ethanol and centrifuged. The obtained product was dried at reduced pressure. The uronic acid content was determined by means of the uronic acid assay according to Blumenkrantz and Abdoe-Hansen (Anal. Biochem., 54

(1973), 484). A calibration curve was generated using polygalacturonic acid (5, 10, 15 and 20 μg). With this calibration curve the uronic acid content in a sample of 20 μg of the product was determined. The obtained result was a content of 95% uronic acid with a yield of 96%.

Partial Oxidation

For partial oxidation, a levan according to the invention (dry weight 2 g, 12.3 mmol) was resuspended in 25 ml water. Next, TEMPO (1% by weight compared to the polysaccharide (0.02 g, 0.13 mmol)) was added, resuspended in 20 min and cooled to 0° C. A solution of hypochlorite (1 ml, 15% solution, 2.1 mmol) was adjusted to pH 9.0 with 3M HCl and cooled down to 0° C. This solution was added to the suspension of the polysaccharide and TEMPO. Within 5 min the mixture became a solid gel.

EXAMPLE 4

Adhesion of *Lactobacillus reuteri* Strains to Caco-2 Cell Lines

The adhesion of *Lactobacillus reuteri* strains to Caco-2 cell lines was determined as described below. Firstly, a bacterial suspension was prepared as follows. *Lactobacillus reuteri* strains LB 121, 35-5, K24 and DSM20016 and *L. rhamnosus* LGG (a well known probiotic strain with good adhering properties) were cultured in MRS broth supplemented with 5 $\mu\text{l/ml}$ of methyl-1,2-[^3H]-thymidine at 37° C. for 18–20 h before the adhesion assays. The cultures were harvested by centrifugation, washed with phosphate buffered saline (PBS) and resuspended in PBS or PBS supplemented with 30 g/l sucrose (see Table 3) to a final density of about 2×10^9 cfu/ml. Prior to the adhesion assay, the cell suspensions in PBS with 30 g/l sucrose were incubated for 1 hour at 37° C., whereas the cell suspensions in PBS were kept on ice for 1 hour. After incubation at 37° C., the suspensions in PBS with sucrose were centrifuged and the cells were washed with and resuspended in PBS to a final density of about 2×10^9 cfu/ml.

Caco-2 cells were cultured as follows. Subcultures of Caco-2 cells (ATCC, code HTB 37, human colon adenocarcinoma), stored as frozen stock cultures in liquid nitrogen were used for the adhesion tests. The Caco-2 cells were grown in culture medium consisting of Dulbecco's modified Eagle medium (DMEM), supplemented with heat-inactivated foetal calf serum (10% v/v), non-essential amino acids (1% v/v), L-glutamine (2 mM) and gentamicin (50 $\mu\text{g/ml}$). About 2,000,000 cells were seeded in 75 cm^2 tissue culture flasks containing culture medium and cultured in a humidified incubator at 37° C. in air containing 5% CO_2 . Near confluent Caco-2 cell cultures were harvested by trypsinisation and resuspended in culture medium. The number of cells was established using a Bürker-Türk counting chamber.

TABLE 3

Incubation of the different <i>Lactobacillus</i> strains prior to the adhesion assays.			
Lactobacillus strain	Extra incubation	Polysaccharide produced	Group
reuteri 121	PBS sucrose, 37° C. for 1 hr	glucan and fructan	As
reuteri 35-5	PBS sucrose, 37° C. for 1 hr	glucan	Bs
reuteri K24	PBS sucrose, 37° C. for 1 hr	none	Cs

TABLE 3-continued

Incubation of the different <i>Lactobacillus</i> strains prior to the adhesion assays.			
Lactobacillus strain	Extra incubation	Polysaccharide produced	Group
reuteri 121	PBS on ice	none	D
reuteri	PBS on ice	none	E
DSM20016*	PBS on ice	none	F

*Type strain of *L. reuteri*

For the following experiments a Caco-2 monolayer transport system was used. Caco-2 cells cultured in a two-compartment transport system are commonly used to study the intestinal, epithelial permeability. In this system the Caco-2 cell differentiates into polarized columnar cells after reaching confluency. The Caco-2 system has been shown to simulate the passive and active transcellular transport of electrolytes, sugars, amino acids and lipophilic compounds (Hillgren et al. 1995, Dulfer et al., 1996, Duizer et al., 1997). Also, a clear correlation between the in vivo absorption and the permeability across the monolayers of Caco-2 cells has been reported (Artursson and Karlsson, 1990). For the present transport studies, Caco-2 cells were seeded on semi-permeable filter inserts (12 wells Transwell plates, Costar) at ca. 100,000 cells per filter (growth area $\pm 1 \text{ cm}^2$ containing 2.5 ml culture medium). The cells on the insert were cultured for 17 to 24 days at 37° C. in a humidified incubator containing 5% CO_2 in air. During this culture period the cells have been subjected to an enterocyte-like differentiation. Gentamycin was eliminated from the culture medium two days prior to the adhesion assays.

The adhesion assay was performed as follows. PBS was used as exposure medium. 25 μl of a bacterial suspension (2×10^9 cfu/ml) were added to 0.5 ml medium. The apical side of the Caco-2 monolayers was incubated with the bacterial suspensions for 1 hour at 37° C. After incubation, remaining fluid was removed and the cells were washed three times with 1 ml PBS. Subsequently, the Caco-2 monolayers were digested overnight with 1 ml 0.1M NaOH, 1% SDS. The lysate was mixed with 10 ml Hionic Fluor scintillation liquid and the radioactivity was measured by liquid scintillation counting using a LKB/Wallac scintillation counter. As a control, the radioactivity of the bacterial suspensions was measured. For each test group, the percentage of bacteria attached to the monolayers was calculated. All adhesion tests were performed in quadruple. In Table 4 the results of the bacterial adhesion test to Caco-2 cell lines are given. From the results can be concluded that the glucans and the fructans contribute to the adherence of *Lactobacillus reuteri* to Caco-2 cell lines. This could indicate that *Lactobacillus reuteri* strains producing EPS possess improved probiotic characteristics or that *Lactobacillus reuteri* and its polysaccharides could function as an excellent symbiotic.

TABLE 4

The results of the bacterial adhesion test to Caco-2 cell lines.		
Group (see Table 1)	0% of bacteria bound to the monolayer	
As	6.5	
Bs	5.7	
Cs	1.8	

TABLE 4-continued

The results of the bacterial adhesion test to Caco-2 cellines.	
Group (see Table 1)	0% of bacteria bound to the monolayer
D	2.3
E	0.9
F	1.3

DESCRIPTION OF THE FIGURES

FIG. 1: The nucleic acid (SEQ ID NO: 4) and deduced amino acid sequences (SEQ ID NOS 27 and 1) of the novel inulosucrase of *Lactobacillus reuteri*. Also encompassed within the figure is the comparison peptide (SEQ ID NO: 28). Furthermore, the designations and orientation (< for 3' to 5' and > for 5' to 3') of the primers and the restriction enzymes used for (inverse) PCR, are shown at the right hand side. Putative start codons (ATG, at positions 41 and 68) and stop codon (TAA, at position 2435) are shown in bold. The positions of the primers used for PCR are shown in bold/underlined. The *NheI* restriction sites (at positions 1154 and 2592) used for inverse PCR are underlined. The primers used and their exact positions in the inulosucrase sequence are shown in table 1. Starting at amino acid 690, the 20 PXX (residues 690-749 of SEQ ID NO: 1) repeats are underlined. At amino acid 755 the LPXTG (SEQ ID NO: 5) motif is underlined.

FIG. 2: Dendrogram of bacterial and plant fructosyltransferases. The horizontal distances are a measure for the

difference at the amino acid sequence level. 10% difference is indicated by the upper bar. Bootstrap values (in percentages) are given at the root of each tree. Fructosyltransferases of Gram positive bacteria are indicated in the lower half of the figure (*B. staerothermophilus* SurB; *B. amyloliquefaciens* SacB; *B. subtilis* SacB; *S. mutans* SacB; *L. reuteri* FtfA (inulosucrase); *S. salivarius* Ftf). Plant fructosyltransferases are indicated in the middle part of the figure (*Cynara scolymus* Ss-1 ftf; *Allium cepa* F-6 gtf; *Hordeum vulgare* Sf-6 ftf). Fructosyltransferases of Gram negative bacteria are shown in the upper part of the figure (*Z. mobilis* LevU; *Z. mobilis* SucE2; *Z. mobilis* SacB; *E. amylovora* Lcs; *A. diazotrophicus* LsdA).

FIG. 3: The N-terminal (SEQ ID NO: 6) and three internal amino acid sequences (SEQ ID NOS 7-9) of the novel levansucrase of *Lactobacillus reuteri*.

FIG. 4: Parts of an alignment of the deduced amino acid sequences of some bacterial fructosyltransferase genes (SEQ ID NOS 29-40). Sequences in bold indicate the consensus sequences used to construct the degenerated primers 5 ftf, 6 ftf and 12 ftf. (*) indicates a position with a fully conserved amino acid residue. (:) indicates a position with a fully conserved 'strong' group: STA, NEQK, NHQK, NDEQ, QHRK, MILV, MILF, HY, FYW. (.) indicates a position with a fully conserved 'weaker' group: CSA, ATV, SAG, STNK, STPA, SGND, SNDEQK, NDEQHK, NEQHRK, FVLIM, HFY. Groups are according to the Pam250 residue weight matrix described by Altschul et al. (1990) J. Mol. Biol. 215, 403-410.

FIG. 5: The strategy used for the isolation of the inulosucrase gene from *Lactobacillus reuteri* 121 chromosomal DNA.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 40

<210> SEQ ID NO 1

<211> LENGTH: 789

<212> TYPE: PRT

<213> ORGANISM: *Lactobacillus reuteri*

<400> SEQUENCE: 1

```

Met Tyr Lys Ser Gly Lys Asn Trp Ala Val Val Thr Leu Ser Thr Ala
 1           5           10           15

Ala Leu Val Phe Gly Ala Thr Thr Val Asn Ala Ser Ala Asp Thr Asn
 20           25           30

Ile Glu Asn Asn Asp Ser Ser Thr Val Gln Val Thr Thr Gly Asp Asn
 35           40           45

Asp Ile Ala Val Lys Ser Val Thr Leu Gly Ser Gly Gln Val Ser Ala
 50           55           60

Ala Ser Asp Thr Thr Ile Arg Thr Ser Ala Asn Ala Asn Ser Ala Ser
 65           70           75           80

Ser Ala Ala Asn Thr Gln Asn Ser Asn Ser Gln Val Ala Ser Ser Ala
 85           90           95

Ala Ile Thr Ser Ser Thr Ser Ser Ala Ala Ser Leu Asn Asn Thr Asp
100           105           110

Ser Lys Ala Ala Gln Glu Asn Thr Asn Thr Ala Lys Asn Asp Asp Thr
115           120           125

Gln Lys Ala Ala Pro Ala Asn Glu Ser Ser Glu Ala Lys Asn Glu Pro

```

-continued

130	135	140
Ala Val Asn Val Asn Asp Ser Ser Ala Ala Lys Asn Asp Asp Gln Gln		
145	150	155 160
Ser Ser Lys Lys Asn Thr Thr Ala Lys Leu Asn Lys Asp Ala Glu Asn		
	165	170 175
Val Val Lys Lys Ala Gly Ile Asp Pro Asn Ser Leu Thr Asp Asp Gln		
	180	185 190
Ile Lys Ala Leu Asn Lys Met Asn Phe Ser Lys Ala Ala Lys Ser Gly		
	195	200 205
Thr Gln Met Thr Tyr Asn Asp Phe Gln Lys Ile Ala Asp Thr Leu Ile		
	210	215 220
Lys Gln Asp Gly Arg Tyr Thr Val Pro Phe Phe Lys Ala Ser Glu Ile		
	225	230 235 240
Lys Asn Met Pro Ala Ala Thr Thr Lys Asp Ala Gln Thr Asn Thr Ile		
	245	250 255
Glu Pro Leu Asp Val Trp Asp Ser Trp Pro Val Gln Asp Val Arg Thr		
	260	265 270
Gly Gln Val Ala Asn Trp Asn Gly Tyr Gln Leu Val Ile Ala Met Met		
	275	280 285
Gly Ile Pro Asn Gln Asn Asp Asn His Ile Tyr Leu Leu Tyr Asn Lys		
	290	295 300
Tyr Gly Asp Asn Glu Leu Ser His Trp Lys Asn Val Gly Pro Ile Phe		
	305	310 315 320
Gly Tyr Asn Ser Thr Ala Val Ser Gln Glu Trp Ser Gly Ser Ala Val		
	325	330 335
Leu Asn Ser Asp Asn Ser Ile Gln Leu Phe Tyr Thr Arg Val Asp Thr		
	340	345 350
Ser Asp Asn Asn Thr Asn His Gln Lys Ile Ala Ser Ala Thr Leu Tyr		
	355	360 365
Leu Thr Asp Asn Asn Gly Asn Val Ser Leu Ala Gln Val Arg Asn Asp		
	370	375 380
Tyr Ile Val Phe Glu Gly Asp Gly Tyr Tyr Tyr Gln Thr Tyr Asp Gln		
	385	390 395 400
Trp Lys Ala Thr Asn Lys Gly Ala Asp Asn Ile Ala Met Arg Asp Ala		
	405	410 415
His Val Ile Glu Asp Gly Asn Gly Asp Arg Tyr Leu Val Phe Glu Ala		
	420	425 430
Ser Thr Gly Leu Glu Asn Tyr Gln Gly Glu Asp Gln Ile Tyr Asn Trp		
	435	440 445
Leu Asn Tyr Gly Gly Asp Asp Ala Phe Asn Ile Lys Ser Leu Phe Arg		
	450	455 460
Ile Leu Ser Asn Asp Asp Ile Lys Ser Arg Ala Thr Trp Ala Asn Ala		
	465	470 475 480
Ala Ile Gly Ile Leu Lys Leu Asn Lys Asp Glu Lys Asn Pro Lys Val		
	485	490 495
Ala Glu Leu Tyr Ser Pro Leu Ile Ser Ala Pro Met Val Ser Asp Glu		
	500	505 510
Ile Glu Arg Pro Asn Val Val Lys Leu Gly Asn Lys Tyr Tyr Leu Phe		
	515	520 525
Ala Ala Thr Arg Leu Asn Arg Gly Ser Asn Asp Asp Ala Trp Met Asn		
	530	535 540
Ala Asn Tyr Ala Val Gly Asp Asn Val Ala Met Val Gly Tyr Val Ala		
	545	550 555 560

-continued

Asp Ser Leu Thr Gly Ser Tyr Lys Pro Leu Asn Asp Ser Gly Val Val
 565 570 575
 Leu Thr Ala Ser Val Pro Ala Asn Trp Arg Thr Ala Thr Tyr Ser Tyr
 580 585 590
 Tyr Ala Val Pro Val Ala Gly Lys Asp Asp Gln Val Leu Val Thr Ser
 595 600 605
 Tyr Met Thr Asn Arg Asn Gly Val Ala Gly Lys Gly Met Asp Ser Thr
 610 615 620
 Trp Ala Pro Ser Phe Leu Gln Ile Asn Pro Asp Asn Thr Thr Thr
 625 630 635 640
 Val Leu Ala Lys Met Thr Asn Gln Gly Asp Trp Ile Trp Asp Asp Ser
 645 650 655
 Ser Glu Asn Leu Asp Met Ile Gly Asp Leu Asp Ser Ala Ala Leu Pro
 660 665 670
 Gly Glu Arg Asp Lys Pro Val Asp Trp Asp Leu Ile Gly Tyr Gly Leu
 675 680 685
 Lys Pro His Asp Pro Ala Thr Pro Asn Asp Pro Glu Thr Pro Thr Thr
 690 695 700
 Pro Glu Thr Pro Glu Thr Pro Asn Thr Pro Lys Thr Pro Lys Thr Pro
 705 710 715 720
 Glu Asn Pro Gly Thr Pro Gln Thr Pro Asn Thr Pro Asn Thr Pro Glu
 725 730 735
 Ile Pro Leu Thr Pro Glu Thr Pro Lys Gln Pro Glu Thr Gln Thr Asn
 740 745 750
 Asn Arg Leu Pro Gln Thr Gly Asn Asn Ala Asn Lys Ala Met Ile Gly
 755 760 765
 Leu Gly Met Gly Thr Leu Leu Ser Met Phe Gly Leu Ala Glu Ile Asn
 770 775 780
 Lys Arg Arg Phe Asn
 785

<210> SEQ ID NO 2
 <211> LENGTH: 2367
 <212> TYPE: DNA
 <213> ORGANISM: Lactobacillus reuteri

<400> SEQUENCE: 2

atgtataaaa gcggtaaaaa ttgggcagtc gttacactct cgactgctgc gctggtattt 60
 ggtgcaacaa ctgtaaatgc atccgcggac acaaatattg aaaacaatga ttcttctact 120
 gtacaagtta caacaggtga taatgatatt gctgttaaaa gtgtgacact tggtagtggt 180
 caagttagtg cagctagtga tacgactatt agaacttctg ctaatgcaaa tagtgcttct 240
 tctgcgcgta atacacaaaa ttctaacagt caagtagcaa gttctgctgc aataacatca 300
 tctacaagtt ccgcagcttc attaaataac acagatagta aagcggctca agaaaatact 360
 aatacagcca aaaatgatga cagcgaacaa gctgcaccag ctaacgaatc ttctgaagct 420
 aaaaatgaac cagctgttaa cgttaatgat tcttcagctg caaaaaatga tgatcaacaa 480
 tccagtaaaa agaatactac cgctaagtta aacaaggatg ctgaaaacgt tgtaaaaaag 540
 gcgggaattg atcctaacag ttttaactgat gaccagatta aagcattaaa taagatgaac 600
 ttctcgaaag ctgcacaaatc tggtagacaa atgacttata atgatttcca aaagattgct 660
 gatacgttaa tcaacaaaga tggtcggtac acagttccat tctttaaagc aagtgaatc 720
 aaaaatatgc ctgccgctac aactaaagat gcacaaacta atactattga acctttagat 780

-continued

gtatgggatt catggccagt tcaagatggt cggacaggac aagttgctaa ttggaatggc	840
tatcaacttg tcatcgcaat gatgggaatt ccaaaccaaa atgataatca tatctatctc	900
ttatataata agtatgggtg taatgaatta agtcattgga agaattgtagg tccaattttt	960
ggctataatt ctaccggcgt ttcacaagaa tggtcaggat cagctgtttt gaacagtgat	1020
aactctatcc aattatttta tacaagggtg gacacgtctg ataacaatac caatcatcaa	1080
aaaattgcta gcgctactct ttatttaact gataataatg gaaatgtatc actcgctcag	1140
gtacgaaatg actatattgt atttgaagggt gatggctatt actaccaaac ttatgatcaa	1200
tggaaagcta ctaacaaggt tgccgataat attgcaatgc gtgatgctca tgtaattgaa	1260
gatggtaatg gtgatcggtg ccttggtttt gaagcaagta ctggttttga aaattatcaa	1320
ggcgaggacc aaatttataa ctggttaaat tatggcggag atgacgcatt taatatcaag	1380
agcttattta gaattctttc caatgatgat attaagagtc gggcaacttg ggctaagtca	1440
gctatcggtg tcctcaaac aaataaggac gaaaagaatc ctaagggtggc agagttatac	1500
tcaccattaa tttctgcacc aatggtaagc gatgaaattg agcgacccaa tgtagttaa	1560
ttaggtaata aatattactt atttgccgct acccgtttaa atcgaggaag taatgatgat	1620
gcttggtatg atgctaatta tgccgttggt gataatgttg caatggctcg atatgttgct	1680
gatatgctaa ctggatctta taagccatta aatgattctg gagtagtctt gactgcttct	1740
gttcttgcaa actggcggac agcaacttat tcatattatg ctgtcccggt tgccggaaaa	1800
gatgaccaag tattagttac ttcatatatg actaatagaa atggagtagc gggtaaagga	1860
atggattcaa ctgggcacc gagtttctta ctacaaatta acccgataa cacaactact	1920
gttttagcta aaatgactaa tcaaggggat tggatttggg atgattcaag cgaatctt	1980
gatatgattg gtgatttaga ctccgctgct ttacctggcg aacgtgataa acctgttgat	2040
tgggacttaa ttggttatgg attaaaaccg catgatcctg ctacacccaa tgatcctgaa	2100
acgccaacta caccagaaac ccctgagaca cctaatactc ccaaacacc aaagactcct	2160
gaaaatcctg ggacacctca aactcctaata acaccttaata ctccggaat tcctttaact	2220
ccagaaacgc ctaagcaacc tgaaccccaa actaataatc gtttgccaca aactggaaat	2280
aatgccaata aagccatgat tggcctaggt atgggaacat tgcttagtat gtttggctct	2340
gcagaaatta acaaacgtcg atttaac	2367

<210> SEQ ID NO 3

<211> LENGTH: 2394

<212> TYPE: DNA

<213> ORGANISM: Lactobacillus reuteri

<400> SEQUENCE: 3

atgctagaac gcaaggaaca taaaaaatg tataaaagcg gtaaaaattg ggcagtcggt	60
acactctcga ctgctgcgct ggtatttggg gcaacaactg taaatgcac cgcggacaca	120
aatattgaaa acaatgatcc ttctactgta caagttacaa caggtgataa tgatattgct	180
gttaaaagtg tgacacttgg tagtggtcaa gttagtgcag ctagtgtac gactattaga	240
acttctgcta atgcaaatag tgcttcttct gccgctaata cacaaaattc taacagtcaa	300
gtagcaagtt ctgctgcaat aacatcatct acaagttccg cagcttcatt aaataacaca	360
gatagtaag cggtctcaaga aaatactaata acagccaaaa atgatgcac gcaaaaagct	420
gcaccagcta acgaatcttc tgaagctaaa aatgaaccag ctgtaaacgt taatgattct	480

-continued

tcagctgcaa	aaaatgatga	tcaacaatcc	agtaaaaaga	atactaccgc	taagttaaac	540
aaggatgctg	aaaacgttgt	aaaaaaggcg	ggaattgata	ctaacagttt	aactgatgac	600
cagattaaag	cattaaataa	gatgaacttc	tcgaaagctg	caaagtctgg	tacacaaatg	660
acttataatg	atttccaaaa	gattgctgat	acgttaetca	aacaagatgg	tcggtacaca	720
gttccattct	ttaaagcaag	tgaatcaaaa	aatatgcctg	ccgtacaca	taaagatgca	780
caactaata	ctattgaacc	tttagatgta	tgggattcat	ggccagtcca	agatgttcgg	840
acaggacaag	ttgctaattg	gaatggctat	caacttgta	tcgcaatgat	gggaattcca	900
aacccaaatg	ataatcatat	ctatctctta	tataataagt	atggtgataa	tgaattaagt	960
cattggaaga	atgtaggtcc	aatttttggc	tataattcta	ccgcgggttc	acaagaatgg	1020
tcaggatcag	ctgttttgaa	cagtataaac	tctatccaat	tattttatgc	aagggtagac	1080
acgtctgata	acaataccaa	tcataaaaa	attgctagcg	ctactcttta	tttaactgat	1140
aataatggaa	atgtatcact	cgctcaggta	cgaatgact	atattgtatt	tgaaggtgat	1200
ggctattact	acaaaactta	tgatcaatgg	aaagctacta	acaaagggtc	cgataatatt	1260
gcaatgcgtg	atgctcatgt	aattgaagat	ggtaatgggt	atcggtacct	tgtttttgaa	1320
gcaagtactg	gtttggaaaa	ttatcaaggc	gaggacccaa	tttataactg	gttaaatatt	1380
ggcggagatg	acgcatttaa	tatcaagagc	ttatttagaa	ttctttccaa	tgatgatatt	1440
aagagtcggg	caacttgggc	taatgcagct	atcggtatcc	tcaaaactaa	taaggacgaa	1500
aagaatccta	agggtggcga	gttatactca	ccattaattt	ctgcaccaat	ggtaagcgat	1560
gaaattgagc	gaccaaaatg	agttaaatta	ggtaataaat	attacttatt	tgcgcctacc	1620
cgtttaaact	gaggaaagta	tgatgatgct	tggatgaatg	ctaattatgc	cgttggtgat	1680
aatgttgcaa	tggtcggata	tggtgctgat	agtctaactg	gatcttataa	gccattaaat	1740
gattctggag	tagtcttgac	tgcttctgtt	cctgcaaaact	ggcggacagc	aacttattca	1800
tattatgctg	tccccgttgc	cggaaaagat	gaccaagtat	tagttacttc	atatatgact	1860
aatagaaatg	gagtagcggg	taaagggaatg	gattcaactt	gggcaccgag	tttcttacta	1920
caaatcaacc	cggataaac	aactactgtt	ttagctaaaa	tgactaatca	aggggattgg	1980
atttgggatg	attcaagcga	aaatcttgat	atgattgggt	atttagactc	cgctgcttta	2040
cctggcgaac	gtgataaacc	tggtgattgg	gacttaattg	gttatggatt	aaaaccgcac	2100
gatactgcta	caccnaatga	tcctgaacg	ccaactacac	cagaaccccc	tgagacacct	2160
aatactccca	aaacacccaa	gactcctgaa	aatcctggga	cacctcaaac	tcctaataca	2220
cctaatactc	cggaaattcc	tttaactcca	gaaacgccta	agcaacctga	aacccaaact	2280
aataatcggt	tgccacaaac	tggaaataat	gccaataaag	ccatgattgg	cctaggtatg	2340
ggaacattgc	ttagtatgtt	tggtcttgca	gaaattaaca	aacgtcgatt	taac	2394

<210> SEQ ID NO 4
 <211> LENGTH: 2592
 <212> TYPE: DNA
 <213> ORGANISM: Lactobacillus reuteri
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(51)
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (68)..(2434)

<400> SEQUENCE: 4

tac aat ggg gtg gcg gag gtg aag aaa cgg ggt tac ttc tat gct aga 48

-continued

Tyr	Asn	Gly	Val	Ala	Glu	Val	Lys	Lys	Arg	Gly	Tyr	Phe	Tyr	Ala	Arg	
1				5					10					15		
acg	caaggaacat	aaaaaa	atg	tat	aaa	agc	ggg	aaa	aat	tgg	gca	gtc	gtt			100
Thr						Met	Tyr	Lys	Ser	Gly	Lys	Asn	Trp	Ala	Val	Val
								20					25			
aca	ctc	tcg	act	gct	gcg	ctg	gta	ttt	ggg	gca	aca	act	gta	aat	gca	148
Thr	Leu	Ser	Thr	Ala	Ala	Leu	Val	Phe	Gly	Ala	Thr	Thr	Val	Asn	Ala	
	30					35				40						
tcc	gcg	gac	aca	aat	att	gaa	aac	aat	gat	tct	tct	act	gta	caa	gtt	196
Ser	Ala	Asp	Thr	Asn	Ile	Glu	Asn	Asn	Asp	Ser	Ser	Thr	Val	Gln	Val	
	45					50				55					60	
aca	aca	ggg	gat	aat	gat	att	gct	gtt	aaa	agt	gtg	aca	ctt	ggg	agt	244
Thr	Thr	Gly	Asp	Asn	Asp	Ile	Ala	Val	Lys	Ser	Val	Thr	Leu	Gly	Ser	
				65					70					75		
ggg	caa	gtt	agt	gca	gct	agt	gat	acg	act	att	aga	act	tct	gct	aat	292
Gly	Gln	Val	Ser	Ala	Ala	Ser	Asp	Thr	Thr	Ile	Arg	Thr	Ser	Ala	Asn	
				80				85					90			
gca	aat	agt	gct	tct	tct	gcc	gct	aat	aca	caa	aat	tct	aac	agt	caa	340
Ala	Asn	Ser	Ala	Ser	Ser	Ala	Ala	Asn	Thr	Gln	Asn	Ser	Asn	Ser	Gln	
	95					100						105				
gta	gca	agt	tct	gct	gca	ata	aca	tca	tct	aca	agt	tcc	gca	gct	tca	388
Val	Ala	Ser	Ser	Ala	Ala	Ile	Thr	Ser	Ser	Thr	Ser	Ser	Ala	Ala	Ser	
	110					115						120				
tta	aat	aac	aca	gat	agt	aaa	gcg	gct	caa	gaa	aat	act	aat	aca	gcc	436
Leu	Asn	Asn	Thr	Asp	Ser	Lys	Ala	Ala	Gln	Glu	Asn	Thr	Asn	Thr	Ala	
	125					130				135					140	
aaa	aat	gat	gac	acg	caa	aaa	gct	gca	cca	gct	aac	gaa	tct	tct	gaa	484
Lys	Asn	Asp	Asp	Thr	Gln	Lys	Ala	Ala	Pro	Ala	Asn	Glu	Ser	Ser	Glu	
				145					150						155	
gct	aaa	aat	gaa	cca	gct	gta	aac	gtt	aat	gat	tct	tca	gct	gca	aaa	532
Ala	Lys	Asn	Glu	Pro	Ala	Val	Asn	Val	Asn	Asp	Ser	Ser	Ala	Ala	Lys	
				160				165					170			
aat	gat	gat	caa	caa	tcc	agt	aaa	aag	aat	act	acc	gct	aag	tta	aac	580
Asn	Asp	Asp	Gln	Gln	Ser	Ser	Lys	Lys	Asn	Thr	Thr	Ala	Lys	Leu	Asn	
				175				180					185			
aag	gat	gct	gaa	aac	gtt	gta	aaa	aag	gcg	gga	att	gat	cct	aac	agt	628
Lys	Asp	Ala	Glu	Asn	Val	Val	Lys	Lys	Ala	Gly	Ile	Asp	Pro	Asn	Ser	
				190			195					200				
tta	act	gat	gac	cag	att	aaa	gca	tta	aat	aag	atg	aac	ttc	tcg	aaa	676
Leu	Thr	Asp	Asp	Gln	Ile	Lys	Ala	Leu	Asn	Lys	Met	Asn	Phe	Ser	Lys	
	205					210				215					220	
gct	gca	aag	tct	ggg	aca	caa	atg	act	tat	aat	gat	ttc	caa	aag	att	724
Ala	Ala	Lys	Ser	Gly	Thr	Gln	Met	Thr	Tyr	Asn	Asp	Phe	Gln	Lys	Ile	
				225					230					235		
gct	gat	acg	tta	atc	aaa	caa	gat	ggg	cgg	tac	aca	gtt	cca	ttc	ttt	772
Ala	Asp	Thr	Leu	Ile	Lys	Gln	Asp	Gly	Arg	Tyr	Thr	Val	Pro	Phe	Phe	
				240				245					250			
aaa	gca	agt	gaa	atc	aaa	aat	atg	cct	gcc	gct	aca	act	aaa	gat	gca	820
Lys	Ala	Ser	Glu	Ile	Lys	Asn	Met	Pro	Ala	Ala	Thr	Thr	Lys	Asp	Ala	
				255			260						265			
caa	act	aat	act	att	gaa	cct	tta	gat	gta	tgg	gat	tca	tgg	cca	gtt	868
Gln	Thr	Asn	Thr	Ile	Glu	Pro	Leu	Asp	Val	Trp	Asp	Ser	Trp	Pro	Val	
				270			275					280				
caa	gat	gtt	cgg	aca	gga	caa	gtt	gct	aat	tgg	aat	ggc	tat	caa	ctt	916
Gln	Asp	Val	Arg	Thr	Gly	Gln	Val	Ala	Asn	Trp	Asn	Gly	Tyr	Gln	Leu	
	285					290				295					300	
gtc	atc	gca	atg	atg	gga	att	cca	aac	caa	aat	gat	aat	cat	atc	tat	964
Val	Ile	Ala	Met	Met	Gly	Ile	Pro	Asn	Gln	Asn	Asp	Asn	His	Ile	Tyr	
				305					310					315		

-continued

ctc tta tat aat aag tat ggt gat aat gaa tta agt cat tgg aag aat	1012
Leu Leu Tyr Asn Lys Tyr Gly Asp Asn Glu Leu Ser His Trp Lys Asn	
320 325 330	
gta ggt cca att ttt ggc tat aat tct acc gcg gtt tca caa gaa tgg	1060
Val Gly Pro Ile Phe Gly Tyr Asn Ser Thr Ala Val Ser Gln Glu Trp	
335 340 345	
tca gga tca gct gtt ttg aac agt gat aac tct atc caa tta ttt tat	1108
Ser Gly Ser Ala Val Leu Asn Ser Asp Asn Ser Ile Gln Leu Phe Tyr	
350 355 360	
aca agg gta gac acg tct gat aac aat acc aat cat caa aaa att gct	1156
Thr Arg Val Asp Thr Ser Asp Asn Asn Thr Asn His Gln Lys Ile Ala	
365 370 375 380	
agc gct act ctt tat tta act gat aat aat gga aat gta tca ctc gct	1204
Ser Ala Thr Leu Tyr Leu Thr Asp Asn Asn Gly Asn Val Ser Leu Ala	
385 390 395	
cag gta cga aat gac tat att gta ttt gaa ggt gat ggc tat tac tac	1252
Gln Val Arg Asn Asp Tyr Ile Val Phe Glu Gly Asp Gly Tyr Tyr Tyr	
400 405 410	
caa act tat gat caa tgg aaa gct act aac aaa ggt gcc gat aat att	1300
Gln Thr Tyr Asp Gln Trp Lys Ala Thr Asn Lys Gly Ala Asp Asn Ile	
415 420 425	
gca atg cgt gat gct cat gta att gaa gat ggt aat ggt gat cgg tac	1348
Ala Met Arg Asp Ala His Val Ile Glu Asp Gly Asn Gly Asp Arg Tyr	
430 435 440	
ctt gtt ttt gaa gca agt act ggt ttg gaa aat tat caa ggc gag gac	1396
Leu Val Phe Glu Ala Ser Thr Gly Leu Glu Asn Tyr Gln Gly Glu Asp	
445 450 455 460	
caa att tat aac tgg tta aat tat ggc gga gat gac gca ttt aat atc	1444
Gln Ile Tyr Asn Trp Leu Asn Tyr Gly Gly Asp Asp Ala Phe Asn Ile	
465 470 475	
aag agc tta ttt aga att ctt tcc aat gat gat att aag agt cgg gca	1492
Lys Ser Leu Phe Arg Ile Leu Ser Asn Asp Asp Ile Lys Ser Arg Ala	
480 485 490	
act tgg gct aat gca gct atc ggt atc ctc aaa cta aat aag gac gaa	1540
Thr Trp Ala Asn Ala Ala Ile Gly Ile Leu Lys Leu Asn Lys Asp Glu	
495 500 505	
aag aat cct aag gtg gca gag tta tac tca cca tta att tct gca cca	1588
Lys Asn Pro Lys Val Ala Glu Leu Tyr Ser Pro Leu Ile Ser Ala Pro	
510 515 520	
atg gta agc gat gaa att gag cga cca aat gta gtt aaa tta ggt aat	1636
Met Val Ser Asp Glu Ile Glu Arg Pro Asn Val Val Lys Leu Gly Asn	
525 530 535 540	
aaa tat tac tta ttt gcc gct acc cgt tta aat cga gga agt aat gat	1684
Lys Tyr Tyr Leu Phe Ala Ala Thr Arg Leu Asn Arg Gly Ser Asn Asp	
545 550 555	
gat gct tgg atg aat gct aat tat gcc gtt ggt gat aat gtt gca atg	1732
Asp Ala Trp Met Asn Ala Asn Tyr Ala Val Gly Asp Asn Val Ala Met	
560 565 570	
gtc gga tat gtt gct gat agt cta act gga tct tat aag cca tta aat	1780
Val Gly Tyr Val Ala Asp Ser Leu Thr Gly Ser Tyr Lys Pro Leu Asn	
575 580 585	
gat tct gga gta gtc ttg act gct tct gtt cct gca aac tgg cgg aca	1828
Asp Ser Gly Val Val Leu Thr Ala Ser Val Pro Ala Asn Trp Arg Thr	
590 595 600	
gca act tat tca tat tat gct gtc ccc gtt gcc gga aaa gat gac caa	1876
Ala Thr Tyr Ser Tyr Tyr Ala Val Pro Val Ala Gly Lys Asp Asp Gln	
605 610 615 620	
gta tta gtt act tca tat atg act aat aga aat gga gta gcg ggt aaa	1924
Val Leu Val Thr Ser Tyr Met Thr Asn Arg Asn Gly Val Ala Gly Lys	
625 630 635	

-continued

gga atg gat tca act tgg gca ccg agt ttc tta cta caa att aac ccg Gly Met Asp Ser Thr Trp Ala Pro Ser Phe Leu Leu Gln Ile Asn Pro 640 645 650	1972
gat aac aca act act gtt tta gct aaa atg act aat caa ggg gat tgg Asp Asn Thr Thr Thr Val Leu Ala Lys Met Thr Asn Gln Gly Asp Trp 655 660 665	2020
att tgg gat gat tca agc gaa aat ctt gat atg att ggt gat tta gac Ile Trp Asp Asp Ser Ser Glu Asn Leu Asp Met Ile Gly Asp Leu Asp 670 675 680	2068
tcc gct gct tta cct ggc gaa cgt gat aaa cct gtt gat tgg gac tta Ser Ala Ala Leu Pro Gly Glu Arg Asp Lys Pro Val Asp Trp Asp Leu 685 690 695 700	2116
att ggt tat gga tta aaa ccg cat gat cct gct aca cca aat gat cct Ile Gly Tyr Gly Leu Lys Pro His Asp Pro Ala Thr Pro Asn Asp Pro 705 710 715	2164
gaa acg cca act aca cca gaa acc cct gag aca cct aat act ccc aaa Glu Thr Pro Thr Thr Pro Glu Thr Pro Glu Thr Pro Asn Thr Pro Lys 720 725 730	2212
aca cca aag act cct gaa aat cct ggg aca cct caa act cct aat aca Thr Pro Lys Thr Pro Glu Asn Pro Gly Thr Pro Gln Thr Pro Asn Thr 735 740 745	2260
cct aat act ccg gaa att cct tta act cca gaa acg cct aag caa cct Pro Asn Thr Pro Glu Ile Pro Leu Thr Pro Glu Thr Pro Lys Gln Pro 750 755 760	2308
gaa acc caa act aat aat cgt ttg cca caa act gga aat aat gcc aat Glu Thr Gln Thr Asn Asn Arg Leu Pro Gln Thr Gly Asn Asn Ala Asn 765 770 775 780	2356
aaa gcc atg att ggc cta ggt atg gga aca ttg ctt agt atg ttt ggt Lys Ala Met Ile Gly Leu Gly Met Gly Thr Leu Leu Ser Met Phe Gly 785 790 795	2404
ctt gca gaa att aac aaa cgt cga ttt aac taatactttt aaaataaaac Leu Ala Glu Ile Asn Lys Arg Arg Phe Asn 800 805	2454
cgctaagcct taaattcagc ttaacggttt tttattttta aagtttttat tgtaaaaaag	2514
cgaattatca ttaatactaa tgcaattggt gtaagacctt acgacagtag taacaatgaa	2574
tttgcccatc ttgtcgg	2592

<210> SEQ ID NO 5
 <211> LENGTH: 5
 <212> TYPE: PRT
 <213> ORGANISM: Lactobacillus reuteri
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (3)
 <223> OTHER INFORMATION: Any amino acid

<400> SEQUENCE: 5

Leu Pro Xaa Thr Gly
1 5

<210> SEQ ID NO 6
 <211> LENGTH: 27
 <212> TYPE: PRT
 <213> ORGANISM: Lactobacillus reuteri

<400> SEQUENCE: 6

Ala Gln Val Glu Ser Asn Asn Tyr Asn Gly Val Ala Glu Val Asn Thr
 1 5 10 15
 Glu Arg Gln Ala Asn Gly Gln Ile Gly Val Asp
 20 25

-continued

<210> SEQ ID NO 7
 <211> LENGTH: 16
 <212> TYPE: PRT
 <213> ORGANISM: Lactobacillus reuteri

<400> SEQUENCE: 7

Met Ala His Leu Asp Val Trp Asp Ser Trp Pro Val Gln Asp Pro Val
 1 5 10 15

<210> SEQ ID NO 8
 <211> LENGTH: 9
 <212> TYPE: PRT
 <213> ORGANISM: Lactobacillus reuteri

<400> SEQUENCE: 8

Asn Ala Gly Ser Ile Phe Gly Thr Lys
 1 5

<210> SEQ ID NO 9
 <211> LENGTH: 19
 <212> TYPE: PRT
 <213> ORGANISM: Lactobacillus reuteri

<400> SEQUENCE: 9

Val Glu Glu Val Tyr Ser Pro Lys Val Ser Thr Leu Met Ala Ser Asp
 1 5 10 15

Glu Val Glu

<210> SEQ ID NO 10
 <211> LENGTH: 4634
 <212> TYPE: DNA
 <213> ORGANISM: Lactobacillus reuteri
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1220)..(3598)
 <220> FEATURE:
 <221> NAME/KEY: RBS
 <222> LOCATION: (1205)..(1210)
 <220> FEATURE:
 <221> NAME/KEY: modified_base
 <222> LOCATION: (2702)..(2707)
 <223> OTHER INFORMATION: a, c, t, g, other or unknown
 <220> FEATURE:
 <221> NAME/KEY: modified_base
 <222> LOCATION: (3686)..(3698)
 <223> OTHER INFORMATION: a, c, t, g, other or unknown

<400> SEQUENCE: 10

gttaacaaag acaaaatttt atataattct tcaaaattaaa tttccactg taagaacata 60
 aatgggtacc tgtttgatgg gaataatata tttgtaacta accggccggc acctttttt 120
 aatgtgccta ggatgcataa tggatgtaaa ttactagatg gcggttttta tacattaacc 180
 tcgcaggaga gaaaagaagc aattagtaag gatccatatg cagataaatt tattaggcct 240
 tatttaggtg ctaaaaattt cattcatgga actgctaggt actgtatttg gtaaaaggac 300
 gcaaacccga aagatatcca tcaatcgcca tttatactgg atagaatcaa taaagtagcg 360
 gaattcagat cgcagcaaaa aagtaaagat acacaaaaat atgcaaaacg gcccatgcta 420
 acaacacgac ttgcctatta tagccacgat gtacatacgg atatgctgat agtacctgca 480
 acatcatcgc aacgtagaga atatcttcca attggatatg tttcagaaaa gaatattgtg 540
 tcttattcac taatgctaatt ccccaatgct agtaatttta atttcggtat tctagaatct 600
 aaagttcact atatttggtt aaaaaacttt tgcggtcggg tgaagtcgga ttatcggtat 660

-continued

tcaaacacta ttatttataa taattttccct tggccgactg ttggtgacaa gccaggamca	720
acaccatctc tgacactcgc tcaaggtata ttaaatactc gcaagctcta tccagacagc	780
tcactggctg atctttatga tccactaaca atgccragtt gaactcgtaa agctcatgaa	840
gccaatgata aagctgttct taaagcatat ggattgagcc ctaaagctac tgagcaagaa	900
atcgtagaac atctatttaa gatgtatgaa aaactgacta aaggtgaaag ataactttgt	960
aaaaccaata ttttataaag acagtaaatg ttaatttgat aaaaacatat atttaataaa	1020
caaaagtgat ataatacagt agttctttgt attacaaaat acatttaata tctctcagca	1080
ttttgcatac tgggagattt tttattgaca aattgtttga aagtgttat gatgaaaccg	1140
tgtagaaact aattcaattt gataaacggt agacatttct gaggaggaag tcattttgga	1200
gtacaaagaa cataagaaa atg tat aaa gtc ggc aag aat tgg gcc gtt gct	1252
Met Tyr Lys Val Gly Lys Asn Trp Ala Val Ala	
1 5 10	
aca ttg gta tca gct tca att tta atg gga ggg gtt gta acc gct cat	1300
Thr Leu Val Ser Ala Ser Ile Leu Met Gly Gly Val Val Thr Ala His	
15 20 25	
gct gat caa gta gaa agt aac aat tac aac ggt gtt gct gaa gtt aat	1348
Ala Asp Gln Val Glu Ser Asn Asn Tyr Asn Gly Val Ala Glu Val Asn	
30 35 40	
act gaa cgt caa gct aat ggt caa att ggc gta gat gga aaa att att	1396
Thr Glu Arg Gln Ala Asn Gly Gln Ile Gly Val Asp Gly Lys Ile Ile	
45 50 55	
agt gct aac agt aat aca acc agt ggc tcg aca aat caa gaa tca tct	1444
Ser Ala Asn Ser Asn Thr Thr Ser Gly Ser Thr Asn Gln Glu Ser Ser	
60 65 70 75	
gct act aac aat act gaa aat gct gtt gtt aat gaa agc aaa aat act	1492
Ala Thr Asn Asn Thr Glu Asn Ala Val Val Asn Glu Ser Lys Asn Thr	
80 85 90	
aac aat act gaa aat gct gtt gtt aat gaa aac aaa aat act aac aat	1540
Asn Asn Thr Glu Asn Ala Val Val Asn Glu Asn Lys Asn Thr Asn Asn	
95 100 105	
act gaa aat gct gtt gtt aat gaa aac aaa aat act aac aac aca gaa	1588
Thr Glu Asn Ala Val Val Asn Glu Asn Lys Asn Thr Asn Asn Thr Glu	
110 115 120	
aac gat aat agt caa tta aag tta act aat aat gaa caa cca tca gcc	1636
Asn Asp Asn Ser Gln Leu Lys Leu Thr Asn Asn Glu Gln Pro Ser Ala	
125 130 135	
gct act caa gca aac ttg aag aag cta aat cct caa gct gct aag gct	1684
Ala Thr Gln Ala Asn Leu Lys Lys Leu Asn Pro Gln Ala Ala Lys Ala	
140 145 150 155	
ggt caa aat gcc aag att gat gcc ggt agt tta aca gat gat caa att	1732
Val Gln Asn Ala Lys Ile Asp Ala Gly Ser Leu Thr Asp Asp Gln Ile	
160 165 170	
aat gaa tta aat aag att aac ttc tct aag tct gct gaa aag ggt gca	1780
Asn Glu Leu Asn Lys Ile Asn Phe Ser Lys Ser Ala Glu Lys Gly Ala	
175 180 185	
aaa ttg acc ttt aag gac tta gag ggg att ggt aat gct att gtt aag	1828
Lys Leu Thr Phe Lys Asp Leu Glu Gly Ile Gly Asn Ala Ile Val Lys	
190 195 200	
caa gat cca caa tat gct att cct tat tct aat gct aag gaa atc aag	1876
Gln Asp Pro Gln Tyr Ala Ile Pro Tyr Ser Asn Ala Lys Glu Ile Lys	
205 210 215	
aat atg cct gca aca tac act gta gat gcc caa aca ggt aag atg gct	1924
Asn Met Pro Ala Thr Tyr Thr Val Asp Ala Gln Thr Gly Lys Met Ala	
220 225 230 235	

-continued

cat ctt gat gtc tgg gac tct tgg cca gta caa gat cct gtc aca ggt His Leu Asp Val Trp Asp Ser Trp Pro Val Gln Asp Pro Val Thr Gly 240 245 250	1972
tat gta tct aat tac atg ggt tat caa cta gtt att gct atg atg ggt Tyr Val Ser Asn Tyr Met Gly Tyr Gln Leu Val Ile Ala Met Met Gly 255 260 265	2020
att cca aat tcg cca act gga gat aat cat atc tat ctt ctt tac aac Ile Pro Asn Ser Pro Thr Gly Asp Asn His Ile Tyr Leu Leu Tyr Asn 270 275 280	2068
aag tat ggt gat aat gac ttt tct cat tgg cgc aat gca ggt tca atc Lys Tyr Gly Asp Asn Asp Phe Ser His Trp Arg Asn Ala Gly Ser Ile 285 290 295	2116
ttt gga act aaa gaa aca aat gtg ttc caa gaa tgg tca ggt tca gct Phe Gly Thr Lys Glu Thr Asn Val Phe Gln Glu Trp Ser Gly Ser Ala 300 305 310 315	2164
att gta aat gat gat ggt aca att caa cta ttt ttc acc tca aat gat Ile Val Asn Asp Asp Gly Thr Ile Gln Leu Phe Phe Thr Ser Asn Asp 320 325 330	2212
acg tct gat tac aag ttg aat gat caa cgc ctt gct acc gca aca tta Thr Ser Asp Tyr Lys Leu Asn Asp Gln Arg Leu Ala Thr Ala Thr Leu 335 340 345	2260
aac ctt aat gtt gat gat aac ggt gtt tca atc aag agt gtt gat aat Asn Leu Asn Val Asp Asp Asn Gly Val Ser Ile Lys Ser Val Asp Asn 350 355 360	2308
tat caa gtt ttg ttt gaa ggt gat gga ttt cac tac caa act tat gaa Tyr Gln Val Leu Phe Glu Gly Asp Gly Phe His Tyr Gln Thr Tyr Glu 365 370 375	2356
caa ttc gca aac ggc aaa gat cgt gaa aat gat gat tac tgc tta cgt Gln Phe Ala Asn Gly Lys Asp Arg Glu Asn Asp Asp Tyr Cys Leu Arg 380 385 390 395	2404
gac cca cac gtt gtt caa tta gaa aat ggt gat cgt tat ctt gta ttc Asp Pro His Val Val Gln Leu Glu Asn Gly Asp Arg Tyr Leu Val Phe 400 405 410	2452
gaa gct aat act ggg aca gaa gat tac caa agt gac gac caa att tat Glu Ala Asn Thr Gly Thr Glu Asp Tyr Gln Ser Asp Asp Gln Ile Tyr 415 420 425	2500
aat tgg gct aac tat ggt ggc gat gat gcc ttc aat att aag agt tcc Asn Trp Ala Asn Tyr Gly Gly Asp Asp Ala Phe Asn Ile Lys Ser Ser 430 435 440	2548
ttc aag ctt ttg aat aat aag aag gat cgt gaa ttg gct ggt tta gct Phe Lys Leu Leu Asn Asn Lys Lys Asp Arg Glu Leu Ala Gly Leu Ala 445 450 455	2596
aat ggt gca ctt ggt atc tta aag ctc act aac aat caa agt aag cca Asn Gly Ala Leu Gly Ile Leu Lys Leu Thr Asn Asn Gln Ser Lys Pro 460 465 470 475	2644
aag gtt gaa gaa gta tac tca cca ttg gta tct act ttg atg gct tgc Lys Val Glu Glu Val Tyr Ser Pro Leu Val Ser Thr Leu Met Ala Cys 480 485 490	2692
gat gag gta nnn nnn aag ctt ggt gat aag tat tat ctc ttc tcc gta Asp Glu Val Xaa Xaa Lys Leu Gly Asp Lys Tyr Tyr Leu Phe Ser Val 495 500 505	2740
act cgt gta agt cgt ggt tcc gat cgt gaa tta acc gct aag gat aac Thr Arg Val Ser Arg Gly Ser Asp Arg Glu Leu Thr Ala Lys Asp Asn 510 515 520	2788
aca atc gtt ggt gat aac gtt gct atg att ggt tac gtt tcc gat agc Thr Ile Val Gly Asp Asn Val Ala Met Ile Gly Tyr Val Ser Asp Ser 525 530 535	2836
tta atg ggt aag tac aag cca tta aat aac tca ggt gtc gta tta act Leu Met Gly Lys Tyr Lys Pro Leu Asn Asn Ser Gly Val Val Leu Thr 540 545 550 555	2884

-continued

gca tca gta cct gca aac tgg cgt act gct act tat tcc tac tat gca Ala Ser Val Pro Ala Asn Trp Arg Thr Ala Thr Tyr Ser Tyr Tyr Ala 560 565 570	2932
gta cct gta gct ggt cat cct gat caa gta tta att act tct tac atg Val Pro Val Ala Gly His Pro Asp Gln Val Leu Ile Thr Ser Tyr Met 575 580 585	2980
agt aac aag gac ttt gct tca ggt gaa gga aac tat gca act tgg gca Ser Asn Lys Asp Phe Ala Ser Gly Glu Gly Asn Tyr Ala Thr Trp Ala 590 595 600	3028
cca agt ttc tta gta caa atc aat cca gat gac acg aca act gta tta Pro Ser Phe Leu Val Gln Ile Asn Pro Asp Asp Thr Thr Thr Val Leu 605 610 615	3076
gca cgt gca act aac caa ggt gac tgg gtg tgg gac gac tct agt cgg Ala Arg Ala Thr Asn Gln Gly Asp Trp Val Trp Asp Asp Ser Ser Arg 620 625 630 635	3124
aac gat aat atg ctc ggt gtt ctt aaa gaa ggt gca gct aac agt gcc Asn Asp Asn Met Leu Gly Val Leu Lys Glu Gly Ala Ala Asn Ser Ala 640 645 650	3172
gcc tta cca ggt gaa tgg ggt aag cca gtt gac tgg agt ttg att aac Ala Leu Pro Gly Glu Trp Gly Lys Pro Val Asp Trp Ser Leu Ile Asn 655 660 665	3220
aga agt cct ggc tta ggc tta aag cct cat caa cca gtt caa cca aag Arg Ser Pro Gly Leu Gly Leu Lys Pro His Gln Pro Val Gln Pro Lys 670 675 680	3268
att gat caa cct gat caa caa cct tct ggt caa aac act aag aat gtc Ile Asp Gln Pro Asp Gln Gln Pro Ser Gly Gln Asn Thr Lys Asn Val 685 690 695	3316
aca cca ggt aat ggt gat aag cct gct ggt aag gca act cct gat aac Thr Pro Gly Asn Gly Asp Lys Pro Ala Gly Lys Ala Thr Pro Asp Asn 700 705 710 715	3364
act aat att gat cca agt gca caa cct tct ggt caa aac act aat att Thr Asn Ile Asp Pro Ser Ala Gln Pro Ser Gly Gln Asn Thr Asn Ile 720 725 730	3412
gat cca agt gca caa mct tct ggt caa aac act aag aat gtc aca cca Asp Pro Ser Ala Gln Xaa Ser Gly Gln Asn Thr Lys Asn Val Thr Pro 735 740 745	3460
ggt aat gag aaa caa ggt aag aat acc gat gca aaa caa tta cca caa Gly Asn Glu Lys Gln Gly Lys Asn Thr Asp Ala Lys Gln Leu Pro Gln 750 755 760	3508
aca ggt aat aag tct ggt tta gca gga ctt tac gct ggt tca tta ctt Thr Gly Asn Lys Ser Gly Leu Ala Gly Leu Tyr Ala Gly Ser Leu Leu 765 770 775	3556
gcc ttg ttt gga ttg gca gca att gaa aag cgt cac gct taa Ala Leu Phe Gly Leu Ala Ala Ile Glu Lys Arg His Ala 780 785 790	3598
tagagtaaaa aaacatcctc cactcaagtt acaagtagga taatatgtat tattttctacg	3658
cytagtcaag aggrattact ggacatannn nnnnnnnnnn tccagttacc aagtggcata	3718
tagtattatt ccacgctagt caggaggatt actgacatta ttggctacat ggccggtagt	3778
cctcttttct tttgtgacga attgtcaaac caagtgaac ggtttctcaa aaaacacctc	3838
atatgggggt tcataattta acacttttcg aggacggcgg ttcagctgat gttggcagaa	3898
actgacgtcc ttatctgtat aatcatcaat attagccctt ttaggaaagt attcccta	3958
tagscattg gtattttcat tgggtcctct ttcctctggt gaatagggat ctggccaata	4018
gatagctact cctaaacgtc ctgcaatata attcaagcca agaaattcac gcccatgatc	4078
tggagtcaat gaatggacaa attcttttagg aatagacct aagagatcaa ttaagccctg	4138

-continued

```

atatttgaat tcggagaagg ggagttgtcc aacaattgcc gttataatac cagggttaat 4198
acggccctgg gcctctacgg taatattgta tttttggctc agatcagtga tagaaaccca 4258
cagatttagc ttgccggtgg agtgctgctt gaagtcttca attacttcgt taccatgttt 4318
gattgctaata ctgatgtgtc gttgttggtg ttagtagaggc atcataccac ctccctcataa 4378
aataaggatat aacaggaatt tctgtacta tatgatcctt ccaatataat aatattaggc 4438
cgataagaaa tgaccagcta ccatttcttg atgcttagtg aatataatcg gatgatcgt 4498
caccctcaa caatccaatt tcacggaggt gagtaatcat gccgagagct aggaatgatt 4558
ggaggaacga acacgggtcca tgcggcagtg gctatttggg ttttagccaa agcagcgta 4618
ctgcttgcaa aagctt 4634

```

```

<210> SEQ ID NO 11
<211> LENGTH: 792
<212> TYPE: PRT
<213> ORGANISM: Lactobacillus reuteri
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (495)..(496)
<223> OTHER INFORMATION: Any amino acid
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (737)
<223> OTHER INFORMATION: Thr or Pro

```

```

<400> SEQUENCE: 11

```

```

Met Tyr Lys Val Gly Lys Asn Trp Ala Val Ala Thr Leu Val Ser Ala
 1          5          10          15
Ser Ile Leu Met Gly Gly Val Val Thr Ala His Ala Asp Gln Val Glu
 20          25          30
Ser Asn Asn Tyr Asn Gly Val Ala Glu Val Asn Thr Glu Arg Gln Ala
 35          40          45
Asn Gly Gln Ile Gly Val Asp Gly Lys Ile Ile Ser Ala Asn Ser Asn
 50          55          60
Thr Thr Ser Gly Ser Thr Asn Gln Glu Ser Ser Ala Thr Asn Asn Thr
 65          70          75          80
Glu Asn Ala Val Val Asn Glu Ser Lys Asn Thr Asn Asn Thr Glu Asn
 85          90          95
Ala Val Val Asn Glu Asn Lys Asn Thr Asn Asn Thr Glu Asn Ala Val
100          105          110
Val Asn Glu Asn Lys Asn Thr Asn Asn Thr Glu Asn Asp Asn Ser Gln
115          120          125
Leu Lys Leu Thr Asn Asn Glu Gln Pro Ser Ala Ala Thr Gln Ala Asn
130          135          140
Leu Lys Lys Leu Asn Pro Gln Ala Ala Lys Ala Val Gln Asn Ala Lys
145          150          155          160
Ile Asp Ala Gly Ser Leu Thr Asp Asp Gln Ile Asn Glu Leu Asn Lys
165          170          175
Ile Asn Phe Ser Lys Ser Ala Glu Lys Gly Ala Lys Leu Thr Phe Lys
180          185          190
Asp Leu Glu Gly Ile Gly Asn Ala Ile Val Lys Gln Asp Pro Gln Tyr
195          200          205
Ala Ile Pro Tyr Ser Asn Ala Lys Glu Ile Lys Asn Met Pro Ala Thr
210          215          220
Tyr Thr Val Asp Ala Gln Thr Gly Lys Met Ala His Leu Asp Val Trp
225          230          235          240

```

-continued

Asp	Ser	Trp	Pro	Val	Gln	Asp	Pro	Val	Thr	Gly	Tyr	Val	Ser	Asn	Tyr	245	250	255
Met	Gly	Tyr	Gln	Leu	Val	Ile	Ala	Met	Met	Gly	Ile	Pro	Asn	Ser	Pro	260	265	270
Thr	Gly	Asp	Asn	His	Ile	Tyr	Leu	Leu	Tyr	Asn	Lys	Tyr	Gly	Asp	Asn	275	280	285
Asp	Phe	Ser	His	Trp	Arg	Asn	Ala	Gly	Ser	Ile	Phe	Gly	Thr	Lys	Glu	290	295	300
Thr	Asn	Val	Phe	Gln	Glu	Trp	Ser	Gly	Ser	Ala	Ile	Val	Asn	Asp	Asp	305	310	315
Gly	Thr	Ile	Gln	Leu	Phe	Phe	Thr	Ser	Asn	Asp	Thr	Ser	Asp	Tyr	Lys	325	330	335
Leu	Asn	Asp	Gln	Arg	Leu	Ala	Thr	Ala	Thr	Leu	Asn	Leu	Asn	Val	Asp	340	345	350
Asp	Asn	Gly	Val	Ser	Ile	Lys	Ser	Val	Asp	Asn	Tyr	Gln	Val	Leu	Phe	355	360	365
Glu	Gly	Asp	Gly	Phe	His	Tyr	Gln	Thr	Tyr	Glu	Gln	Phe	Ala	Asn	Gly	370	375	380
Lys	Asp	Arg	Glu	Asn	Asp	Asp	Tyr	Cys	Leu	Arg	Asp	Pro	His	Val	Val	385	390	395
Gln	Leu	Glu	Asn	Gly	Asp	Arg	Tyr	Leu	Val	Phe	Glu	Ala	Asn	Thr	Gly	405	410	415
Thr	Glu	Asp	Tyr	Gln	Ser	Asp	Asp	Gln	Ile	Tyr	Asn	Trp	Ala	Asn	Tyr	420	425	430
Gly	Gly	Asp	Asp	Ala	Phe	Asn	Ile	Lys	Ser	Ser	Phe	Lys	Leu	Leu	Asn	435	440	445
Asn	Lys	Lys	Asp	Arg	Glu	Leu	Ala	Gly	Leu	Ala	Asn	Gly	Ala	Leu	Gly	450	455	460
Ile	Leu	Lys	Leu	Thr	Asn	Asn	Gln	Ser	Lys	Pro	Lys	Val	Glu	Glu	Val	465	470	475
Tyr	Ser	Pro	Leu	Val	Ser	Thr	Leu	Met	Ala	Cys	Asp	Glu	Val	Xaa	Xaa	485	490	495
Lys	Leu	Gly	Asp	Lys	Tyr	Tyr	Leu	Phe	Ser	Val	Thr	Arg	Val	Ser	Arg	500	505	510
Gly	Ser	Asp	Arg	Glu	Leu	Thr	Ala	Lys	Asp	Asn	Thr	Ile	Val	Gly	Asp	515	520	525
Asn	Val	Ala	Met	Ile	Gly	Tyr	Val	Ser	Asp	Ser	Leu	Met	Gly	Lys	Tyr	530	535	540
Lys	Pro	Leu	Asn	Asn	Ser	Gly	Val	Val	Leu	Thr	Ala	Ser	Val	Pro	Ala	545	550	555
Asn	Trp	Arg	Thr	Ala	Thr	Tyr	Ser	Tyr	Tyr	Ala	Val	Pro	Val	Ala	Gly	565	570	575
His	Pro	Asp	Gln	Val	Leu	Ile	Thr	Ser	Tyr	Met	Ser	Asn	Lys	Asp	Phe	580	585	590
Ala	Ser	Gly	Glu	Gly	Asn	Tyr	Ala	Thr	Trp	Ala	Pro	Ser	Phe	Leu	Val	595	600	605
Gln	Ile	Asn	Pro	Asp	Asp	Thr	Thr	Thr	Val	Leu	Ala	Arg	Ala	Thr	Asn	610	615	620
Gln	Gly	Asp	Trp	Val	Trp	Asp	Asp	Ser	Ser	Arg	Asn	Asp	Asn	Met	Leu	625	630	635
Gly	Val	Leu	Lys	Glu	Gly	Ala	Ala	Asn	Ser	Ala	Ala	Leu	Pro	Gly	Glu	645	650	655
Trp	Gly	Lys	Pro	Val	Asp	Trp	Ser	Leu	Ile	Asn	Arg	Ser	Pro	Gly	Leu			

-continued

660	665	670
Gly Leu Lys Pro His Gln Pro Val Gln Pro Lys Ile Asp Gln Pro Asp 675 680 685		
Gln Gln Pro Ser Gly Gln Asn Thr Lys Asn Val Thr Pro Gly Asn Gly 690 695 700		
Asp Lys Pro Ala Gly Lys Ala Thr Pro Asp Asn Thr Asn Ile Asp Pro 705 710 715 720		
Ser Ala Gln Pro Ser Gly Gln Asn Thr Asn Ile Asp Pro Ser Ala Gln 725 730 735		
Xaa Ser Gly Gln Asn Thr Lys Asn Val Thr Pro Gly Asn Glu Lys Gln 740 745 750		
Gly Lys Asn Thr Asp Ala Lys Gln Leu Pro Gln Thr Gly Asn Lys Ser 755 760 765		
Gly Leu Ala Gly Leu Tyr Ala Gly Ser Leu Leu Ala Leu Phe Gly Leu 770 775 780		
Ala Ala Ile Glu Lys Arg His Ala 785 790		
 <210> SEQ ID NO 12 <211> LENGTH: 24 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Description of Artificial Sequence: Primer <400> SEQUENCE: 12 ctgataataa tggaaatgta tcac		
		24
 <210> SEQ ID NO 13 <211> LENGTH: 26 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Description of Artificial Sequence: Primer <400> SEQUENCE: 13 catgatcata agtttggtag taatag		
		26
 <210> SEQ ID NO 14 <211> LENGTH: 24 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Description of Artificial Sequence: Primer <400> SEQUENCE: 14 gtgatacatt tccattatta tcag		
		24
 <210> SEQ ID NO 15 <211> LENGTH: 26 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Description of Artificial Sequence: Primer <400> SEQUENCE: 15 ctattactac caaacttatg atcatg		
		26
 <210> SEQ ID NO 16 <211> LENGTH: 38 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence		

-continued

```

<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 16

ccatggccat ggtagaacgc aaggaacata aaaaaatg
38

<210> SEQ ID NO 17
<211> LENGTH: 38
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 17

agatctagat ctgttaaact gacgtttgtt aatttctg
38

<210> SEQ ID NO 18
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (6)
<223> OTHER INFORMATION: a, c, t, g, other or unknown
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (15)
<223> OTHER INFORMATION: a, c, t, g, other or unknown

<400> SEQUENCE: 18

gaygtntggg aywantgggc c
21

<210> SEQ ID NO 19
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (3)
<223> OTHER INFORMATION: a, c, t, g, other or unknown
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (6)
<223> OTHER INFORMATION: a, c, t, g, other or unknown
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (9)
<223> OTHER INFORMATION: a, c, t, g, other or unknown
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (12)
<223> OTHER INFORMATION: a, c, t, g, other or unknown
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 19

gtngcswnc cswccayts ytg
23

<210> SEQ ID NO 20
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 20

gaatgtaggt ccaatttttg gc
22

```


-continued

<210> SEQ ID NO 21
 <211> LENGTH: 22
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 21

cctgtccgaa catcttgaaac tg

22

<210> SEQ ID NO 22
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Primer
 <220> FEATURE:
 <221> NAME/KEY: modified_base
 <222> LOCATION: (6)
 <223> OTHER INFORMATION: a, c, t, g, other or unknown
 <220> FEATURE:
 <221> NAME/KEY: modified_base
 <222> LOCATION: (9)
 <223> OTHER INFORMATION: a, c, t, g, other or unknown
 <220> FEATURE:
 <221> NAME/KEY: modified_base
 <222> LOCATION: (12)
 <223> OTHER INFORMATION: a, c, t, g, other or unknown
 <220> FEATURE:
 <221> NAME/KEY: modified_base
 <222> LOCATION: (18)
 <223> OTHER INFORMATION: a, c, t, g, other or unknown
 <220> FEATURE:
 <221> NAME/KEY: modified_base
 <222> LOCATION: (21)
 <223> OTHER INFORMATION: a, c, t, g, other or unknown

<400> SEQUENCE: 22

arraanswng gngcvmangt nsw

23

<210> SEQ ID NO 23
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Primer
 <220> FEATURE:
 <221> NAME/KEY: modified_base
 <222> LOCATION: (9)
 <223> OTHER INFORMATION: a, c, t, g, other or unknown
 <220> FEATURE:
 <221> NAME/KEY: modified_base
 <222> LOCATION: (11)
 <223> OTHER INFORMATION: a, c, t, g, other or unknown
 <220> FEATURE:
 <221> NAME/KEY: modified_base
 <222> LOCATION: (15)
 <223> OTHER INFORMATION: a, c, t, g, other or unknown
 <220> FEATURE:
 <221> NAME/KEY: modified_base
 <222> LOCATION: (21)
 <223> OTHER INFORMATION: a, c, t, g, other or unknown

<400> SEQUENCE: 23

tayaayggng tngcngargt naa

23

<210> SEQ ID NO 24
 <211> LENGTH: 22
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Primer

-continued

<400> SEQUENCE: 24

ccgaccatct tgtttgatta ac

22

<210> SEQ ID NO 25

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 25

aaytataayg gygttgcryg aagt

24

<210> SEQ ID NO 26

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<220> FEATURE:

<221> NAME/KEY: modified_base

<222> LOCATION: (9)

<223> OTHER INFORMATION: a, c, t, g, other or unknown

<400> SEQUENCE: 26

taccgnwsnc tacttcaact t

21

<210> SEQ ID NO 27

<211> LENGTH: 17

<212> TYPE: PRT

<213> ORGANISM: Lactobacillus reuteri

<400> SEQUENCE: 27

Tyr	Asn	Gly	Val	Ala	Glu	Val	Lys	Lys	Arg	Gly	Tyr	Phe	Tyr	Ala	Arg
1					5				10					15	

Thr

<210> SEQ ID NO 28

<211> LENGTH: 17

<212> TYPE: PRT

<213> ORGANISM: Lactobacillus reuteri

<400> SEQUENCE: 28

Tyr	Asn	Gly	Val	Ala	Glu	Val	Asn	Thr	Glu	Arg	Gln	Ala	Asn	Gly	Gly
1					5				10					15	

Ile

<210> SEQ ID NO 29

<211> LENGTH: 14

<212> TYPE: PRT

<213> ORGANISM: Bacillus amyloliquefaciens

<400> SEQUENCE: 29

Gly	Leu	Asp	Val	Trp	Asp	Ser	Trp	Pro	Leu	Gln	Asn	Ala	Asp
1				5					10				

<210> SEQ ID NO 30

<211> LENGTH: 14

<212> TYPE: PRT

<213> ORGANISM: Bacillus subtilis

<400> SEQUENCE: 30

Gly Leu Asp Val Trp Asp Ser Trp Pro Leu Gln Asn Ala Asp

-continued

1	5	10
---	---	----

<210> SEQ ID NO 31
 <211> LENGTH: 14
 <212> TYPE: PRT
 <213> ORGANISM: Streptococcus mutans
 <400> SEQUENCE: 31

Asp	Leu	Asp	Val	Trp	Asp	Ser	Trp	Pro	Val	Gln	Asp	Ala	Lys
1			5						10				

<210> SEQ ID NO 32
 <211> LENGTH: 14
 <212> TYPE: PRT
 <213> ORGANISM: Streptococcus salivarius
 <400> SEQUENCE: 32

Glu	Ile	Asp	Val	Trp	Asp	Ser	Trp	Pro	Val	Gln	Asp	Ala	Lys
1			5						10				

<210> SEQ ID NO 33
 <211> LENGTH: 16
 <212> TYPE: PRT
 <213> ORGANISM: Bacillus amyloliquefaciens
 <400> SEQUENCE: 33

Gln	Thr	Gln	Glu	Trp	Ser	Gly	Ser	Ala	Thr	Phe	Thr	Ser	Asp	Gly	Lys
1			5						10					15	

<210> SEQ ID NO 34
 <211> LENGTH: 16
 <212> TYPE: PRT
 <213> ORGANISM: Bacillus subtilis
 <400> SEQUENCE: 34

Gln	Thr	Gln	Glu	Trp	Ser	Gly	Ser	Ala	Thr	Phe	Thr	Ser	Asp	Gly	Lys
1			5						10					15	

<210> SEQ ID NO 35
 <211> LENGTH: 16
 <212> TYPE: PRT
 <213> ORGANISM: Streptococcus mutans
 <400> SEQUENCE: 35

Leu	Thr	Gln	Glu	Trp	Ser	Gly	Ser	Ala	Thr	Val	Asn	Glu	Asp	Gly	Ser
1			5						10					15	

<210> SEQ ID NO 36
 <211> LENGTH: 16
 <212> TYPE: PRT
 <213> ORGANISM: Streptococcus salivarius
 <400> SEQUENCE: 36

Asp	Asp	Gln	Gln	Trp	Ser	Gly	Ser	Ala	Thr	Val	Asn	Ser	Asp	Gly	Ser
1			5						10					15	

<210> SEQ ID NO 37
 <211> LENGTH: 11
 <212> TYPE: PRT
 <213> ORGANISM: Bacillus amyloliquefaciens
 <400> SEQUENCE: 37

Lys	Ala	Thr	Phe	Gly	Pro	Ser	Phe	Leu	Met	Asn
1			5					10		

-continued

<210> SEQ ID NO 38
 <211> LENGTH: 11
 <212> TYPE: PRT
 <213> ORGANISM: *Bacillus subtilis*
 <400> SEQUENCE: 38

Gln Ser Thr Phe Ala Pro Ser Phe Leu Leu Asn
 1 5 10

<210> SEQ ID NO 39
 <211> LENGTH: 11
 <212> TYPE: PRT
 <213> ORGANISM: *Streptococcus mutans*
 <400> SEQUENCE: 39

Asn Ser Thr Trp Ala Pro Ser Phe Leu Ile Gln
 1 5 10

<210> SEQ ID NO 40
 <211> LENGTH: 11
 <212> TYPE: PRT
 <213> ORGANISM: *Streptococcus salivarius*
 <400> SEQUENCE: 40

Lys Ser Thr Trp Ala Pro Ser Phe Leu Ile Lys
 1 5 10

What is claimed is:

1. A process of producing a fructo-oligosaccharide or fructo-polysaccharide, having $\beta(2-1)$ linked D-fructosyl units comprising forming a mixture by combining sucrose with at least one reaction partner selected from the group consisting of:

- a) a protein having fructosyltransferase activity, which exhibits at least 85% amino acid identity, as determined by a BLAST algorithm, with an amino acid sequence of SEQ ID No. 1, and
- b) a recombinant host cell containing one or more copies of a nucleic acid construct encoding for said protein (a) and capable of expressing said protein;

wherein said reaction partner interacts with sucrose to produce a fructo-oligosaccharide or fructo-polysaccharide.

2. The process according to claim 1, wherein said protein is a recombinant protein.

3. A process according to claim 1, further comprising chemically modifying said fructo-oligosaccharide or fructo-polysaccharide by simultaneous 3- and 4-oxidation, 1- or 6-oxidation, phosphorylation, acylation, hydroxyalkylation, carboxymethylation or amino-alkylation of one or more anhydrofructose units, or by hydrolysis.

4. The process according to claim 1, further comprising adding a food or beverage composition to said mixture to obtain a prebiotic composition.

5. The process according to claim 1, further comprising adding to said mixture a *Lactobacillus* strain capable of producing an oligosaccharide or polysaccharide and optionally a food or beverage composition, to obtain a synbiotic composition.

6. A process of producing a fructo-oligosaccharide or fructo-polysaccharide, having $\beta(2-1)$ linked D-fructosyl units comprising combining sucrose and a protein to form a mixture, said protein having fructosyltransferase activity, which exhibits at least 85% amino acid identity, as determined by a BLAST algorithm, with an amino acid sequence of SEQ ID No. 1, and

interacting said sucrose with said protein to produce said fructo-oligosaccharide or fructo-polysaccharide.

7. A process for producing a fructo-oligosaccharide or fructo-polysaccharide, having $\beta(2-6)$ linked D-fructosyl units comprising forming a mixture by combining sucrose with a reaction partner, wherein said reaction partner is a recombinant host cell containing one or more copies of a nucleic acid construct encoding for a protein having fructosyltransferase activity, which exhibits at least 85% amino acid identity, as determined by a BLAST algorithm, with an amino acid sequence of SEQ ID No. 11, and wherein said reaction partner interacts with sucrose to provide a fructo-oligosaccharide or fructo-polysaccharide.

8. A process according to claim 7, further comprising chemically modifying said fructo-oligosaccharide or fructo-polysaccharide by simultaneous 3- and 4-oxidation, 1- or 6-oxidation, phosphorylation, acylation, hydroxyalkylation, carboxymethylation or amino-alkylation of one or more anhydrofructose units, or by hydrolysis.

* * * * *